




Lomefloxacin

Revised: January 18, 2021.

CASRN: 98079-51-7



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of lomefloxacin during breastfeeding. Fluoroquinolones have traditionally not been used in infants because of concern about adverse effects on the infants' developing joints. However, recent studies indicate little risk.[1,2] The calcium in milk might prevent absorption of the small amounts of fluoroquinolones in milk,[3] but insufficient data exist to prove or disprove this assertion. Use of lomefloxacin is acceptable in nursing mothers with monitoring of the infant for possible effects on the gastrointestinal flora, such as diarrhea or candidiasis (thrush, diaper rash). However, it is preferable to use an alternate drug for which safety information is available.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Ciprofloxacin, Levofloxacin

References

1. Gürpınar AN, Balkan E, Kiliç N, et al. The effects of a fluoroquinolone on the growth and development of infants. *J Int Med Res.* 1997;25:302–6. PubMed PMID: 9364293.
2. van den Oever HL, Versteegh FG, Thewessen EA, et al. Ciprofloxacin in preterm neonates: Case report and review of the literature. *Eur J Pediatr.* 1998;157:843–5. PubMed PMID: 9809826.
3. Fleiss PM. The effect of maternal medications on breast-feeding infants. *J Hum Lact* 1992;8:7. Letter. PMID: 1558663

Substance Identification

Substance Name

Lomefloxacin

CAS Registry Number

98079-51-7

Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Antibacterial Agents

Quinolones

Fluoroquinolones