




## Apomorphine

Revised: July 18, 2022.

CASRN: 58-00-4



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of apomorphine during breastfeeding. If apomorphine is required by the mother, it is not a reason to discontinue breastfeeding. However, apomorphine inhibits prolactin release in animals and might interfere with establishment of lactation. An alternate drug may be preferred, especially while nursing a newborn or preterm infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Apomorphine

### CAS Registry Number

58-00-4

### Drug Class

Breastfeeding

Lactation

Milk, Human

Antiparkinson Agents

Dopamine Agents

Dopamine Agonists

Emetics

Gastrointestinal Agents