



Dostarlimab

Revised: May 17, 2021.

CASRN: 2022215-59-2

Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of dostarlimab during breastfeeding. Because dostarlimab is a large protein molecule with a molecular weight of about 144,000 Da, the amount in milk is likely to be very low. It is also likely to be partially destroyed in the infant's gastrointestinal tract and absorption by the infant is probably minimal. Because of the potential for serious adverse reactions in the breastfed infant, the manufacturer recommends that women not breastfeed during treatment and for 4 months after the final dose.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Dostarlimab

CAS Registry Number

2022215-59-2

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Drug Class

Breast Feeding

Lactation

Antibodies, Monoclonal, Humanized

Antineoplastic Agents