




## Pimavanserin

Revised: November 30, 2022.

CASRN: 706779-91-1



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the clinical use of pimavanserin during breastfeeding. If pimavanserin is required by the mother, it is not a reason to discontinue breastfeeding. However, an alternate drug may be preferred, especially while nursing a newborn or preterm infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Haloperidol, Olanzapine, Quetiapine, Risperidone

## Substance Identification

### Substance Name

Pimavanserin

### CAS Registry Number

706779-91-1

### Drug Class

Breast Feeding

Lactation

Milk, Human

Antiparkinson Agents

Antipsychotic Agents

Serotonin 5-HT<sub>2</sub> Receptor Antagonists