

3.5 ATAZANAVIR (ATV)

FORMULATIONS

Tablets	Capsules	Liquid	FDC
None	100 mg 150 mg 200 mg 300 mg	None	None

DOSE AND FREQUENCY OF DOSING

Target dose ATV/RTV

Treatment-naïve

- Weight 15 kg to less than 25 kg: 150 mg ATV/80 mg RTV
- Weight 25 kg to less than 32 kg: 200 mg ATV/100 mg RTV
- Weight 32 kg to less than 39 kg: 250 mg ATV/100 mg RTV

Treatment-experienced

- Weight 25 kg to less than 32 kg: 200 mg ATV/100 mg RTV
- Weight 32 kg to less than 39 kg: 250 mg ATV/100 mg RTV

Maximum dose

- ATV 300 mg/RTV 100 mg once daily

Administration

- Should be taken with food

Storage

- Store capsules at 25°C (range 15 – 30°C).

OTHER COMMENTS

General

- To be used in combination with RTV in paediatric patients
- Recommended for patients aged 6 to <18 years of age
- Not to be used in patients less than 3 months of age due to risk of kernicterus. There are insufficient data for patients less than 6 years of age.
- Dosage is based on body weight (8.5 mg/kg for weights 15 kg to less than 20 kg, and 7 mg/kg for weight 20 kg or more)

Pharmacokinetic data

- Available for children aged 3 months to 21 years

Major drug interactions

- None reported

Ref: http://packageinserts.bms.com/pi/pi_revataz.pdf