

**Table E-8. Study characteristics for trials comparing nonprescription nonhormone with nonprescription nonhormone**

<b>Study</b>	<b>Country</b>	<b>Sites</b>	<b>Analyzed</b>	<b>Design</b>	<b>Study Arms</b>	<b>F/U (weeks)</b>	<b>Industry Funding</b>	<b>Editorial Assistance</b>
Liske 2002	Poland	4	149	RCT Active Compare	2	12		No/NR
Hidalgo 2006	Spain	Multi	925	RCT Active Compare	2	26		No/NR
Zervoudis (a) 2008	Greece	Multi	62	RCT Active Compare	2	52		No/NR
Agosta 2011	Italy	91	636	RCT Active Compare	2	12		No/NR
Le Donne 2011	Italy	1	62	RCT Active Compare	2	13		No/NR
Virojchaiwong 2011	Thailand	1	52	RCT Active Compare	2	26		No/NR
Yang 2012	Taiwan	4	130	RCT Active Compare	2	24		No/NR

(a): data came from a conference abstract; (c): data came from posted results on the clinical trial registry; (d): duplicate patient population with other included article; F/U: followup; (m): trial contains data from multiple publications; Multi: multicenter trial; NR: not reported; Plac: placebo; RCT: randomized controlled trial; (SIP); data came from a