




Butoconazole

Revised: October 31, 2018.

CASRN: 64872-76-0



Drug Levels and Effects

Summary of Use during Lactation

Vaginal butoconazole has not been studied during breastfeeding. About 5.5% of a vaginal dose is absorbed and its plasma half-life is 21 to 24 hours. Because there is no published experience with butoconazole during breastfeeding, other agents may be preferred, especially while nursing a newborn or preterm infant.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Clotrimazole, Miconazole

Substance Identification

Substance Name

Butoconazole

CAS Registry Number

64872-76-0

Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Antifungal Agents

Dermatologic Agents