



U.S. National Library of Medicine  
National Center for Biotechnology Information

**NLM Citation:** Wallace SE, Bean LJH. Resources for Genetics Professionals — Genetic Disorders Caused by Imprinting Errors and Uniparental Disomy Not Detectable by Sequence Analysis. 2017 Mar 14. In: Adam MP, Feldman J, Mirzaa GM, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024.  
**Bookshelf URL:** <https://www.ncbi.nlm.nih.gov/books/>



## Resources for Genetics Professionals – Genetic Disorders Caused by Imprinting Errors and Uniparental Disomy Not Detectable by Sequence Analysis

Stephanie E Wallace, MD<sup>1,2</sup> and Lora JH Bean, PhD<sup>3,4</sup>

Created: March 14, 2017.

### Imprinting

Imprinting is the process by which maternally and paternally derived chromosomes are uniquely chemically modified (usually by methylation), leading to different expression of a certain gene or genes on those chromosomes depending on their parental origin. Patterns of gene expression and repression of gene expression vary between imprinted regions.

Methylation, the attachment of methyl groups to DNA at cytosine bases, is correlated with reduced transcription of a gene and is thought to be the principal mechanism of imprinting. An assay designed to detect the methylation pattern at a specific chromosome locus (e.g., methylation-sensitive multiplex ligation probe analysis [MS-MLPA], methylation-sensitive quantitative PCR [MS-qPCR], Southern blotting) is necessary to identify epigenetic imprinting errors. Sequencing techniques including next-generation sequencing used in exome sequencing and genome sequencing do not identify methylation patterns.

### Uniparental Disomy

Uniparental disomy (UPD) is the situation in which both members of a chromosome pair or segments of a chromosome pair are inherited from one parent and neither is inherited from the other parent; uniparental disomy can result in an abnormal phenotype if genes within the chromosome segment are imprinted.

The ability of sequence analysis to identify UPD by sequencing DNA from the proband alone depends on the type: uniparental isodisomy versus uniparental heterodisomy.

- **Uniparental isodisomy** (i.e., two copies of a single chromosome or chromosome segment are inherited from one parent and no copy is inherited from the other parent) may be identified by long stretches of

**Author Affiliations:** 1 Senior Editor, GeneReviews; Email: editor2@uw.edu. 2 Clinical Professor, Pediatrics, University of Washington, Seattle, Washington; Email: editor2@uw.edu. 3 Molecular Genetics Editor, GeneReviews. 4 Senior Director, Laboratory Quality Assurance, Perkin-Elmer Genomics, Inc, Pittsburgh, Pennsylvania.

Copyright © 1993-2024, University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved.

homozygosity by sequencing techniques; however, further testing (e.g., chromosomal microarray analysis) may be required to distinguish a deletion from uniparental isodisomy.

- **Uniparental heterodisomy** (i.e., both chromosomes of a chromosome pair or chromosome segment are inherited from one parent and no copy is inherited from the other parent) cannot be identified by sequencing techniques unless recombination has reduced regions of the chromosome to homozygosity. Other test methods such as methylation studies or parental testing (e.g., short tandem repeat (STR) analysis or SNP array) are required to identify uniparental heterodisomy.

The table below provides a list of disorders caused by imprinting errors and UPD organized by chromosome locus.

**Table.** Genetic Disorders Caused by Imprinting Errors and Uniparental Disomy <sup>1</sup> Not Detectable by Sequence Analysis

Chromosome Locus	Disorder <sup>2</sup>	Proportion of Disorder Due to Imprinting Error Not Detectable by Sequence Analysis	Proportion of Disorder Due to UPD (Parental Origin of Disomic Chromosome)
6q24.2	<a href="#">Transient neonatal diabetes mellitus, 6q24 related</a>	21%	41% (paternal)
chr 7	<a href="#">Silver-Russell syndrome</a>		7%-10% (maternal)
11p15.5	<a href="#">Silver-Russell syndrome</a>	35%-50%	
11p15.5	<a href="#">Beckwith-Wiedemann syndrome</a>	55%	~20% (paternal)
11p15.5	<a href="#">Isolated hemihyperplasia (OMIM 235000)</a>	6% <sup>3</sup>	16% <sup>3</sup>
11p15.5	<a href="#">Isolated Wilms tumor (See <a href="#">Wilms Tumor Predisposition.</a>)</a>	<1% <sup>4</sup>	<1% <sup>4</sup>
14q32	<a href="#">Temple syndrome (OMIM 616222)</a>		100% (maternal)
14q32	<a href="#">Kagami-Ogata syndrome</a>		100% (paternal)
15q11.2-q13	<a href="#">Prader-Willi syndrome</a>	2%	20%-30% (maternal)
15q11.2-q13	<a href="#">Angelman syndrome</a>	80%	~7% (paternal)
20q13	<a href="#">Pseudohypoparathyroidism 1B (See <a href="#">Disorders of GNAS Inactivation.</a>)</a>	~10%-30%	~10% <sup>5</sup> (paternal)
chr 20	<a href="#">UPD(20)mat <sup>6</sup></a>		12 individuals (maternal)

UPD = uniparental disomy

1. Long stretches of homozygosity caused by uniparental isodisomy may be detected by exome or genome sequence analysis; however, not all mechanisms by which uniparental disomy occurs result in stretches of homozygosity [Kearney et al 2011]. The ability of a laboratory to recognize stretches of homozygosity will vary.

2. For more information see hyperlinked *GeneReview*. An OMIM phenotype entry (if available) is provided if there is no *GeneReview*.

3. Shuman et al [2006], Soellner et al [2017]

4. Scott et al [2008]

5. Takatani et al [2015]

6. Mulchandani et al [2016]

## References

Kearney HM, Kearney JB, Conlin LK. Diagnostic implications of excessive homozygosity detected by SNP-based microarrays: consanguinity, uniparental disomy, and recessive single-gene mutations. *Clin Lab Med.* 2011;31:595–613. PubMed PMID: 22118739.

Mulchandani S, Bhoj EJ, Luo M, Powell-Hamilton N, Jenny K, Gripp KW, Elbracht M, Eggermann T, Turner CL, Temple IK, Mackay DJ, Dubbs H, Stevenson DA, Slattery L, Zackai EH, Spinner NB, Krantz ID, Conlin LK.

Maternal uniparental disomy of chromosome 20: a novel imprinting disorder of growth failure. *Genet Med.* 2016;18:309–15. PubMed PMID: 26248010.

Scott RH, Douglas J, Baskcomb L, Huxter N, Barker K, Hanks S, Craft A, Gerrard M, Kohler JA, Levitt GA, Picton S, Pizer B, Ronghe MD, Williams D., Factors Associated with Childhood Tumours (FACT) Collaboration. Cook JA, Pujol P, Maher ER, Birch JM, Stiller CA, Pritchard-Jones K, Rahman N. Constitutional 11p15 abnormalities, including heritable imprinting center mutations, cause nonsyndromic Wilms tumor. *Nat Genet.* 2008;40:1329–34. PubMed PMID: 18836444.

Shuman C, Smith AC, Steele L, Ray PN, Clericuzio C, Zackai E, Parisi MA, Meadows AT, Kelly T, Tichauer D, Squire JA, Sadowski P, Weksberg R. Constitutional UPD for chromosome 11p15 in individuals with isolated hemihyperplasia is associated with high tumor risk and occurs following assisted reproductive technologies. *Am J Med Genet A.* 2006;140:1497–503. PubMed PMID: 16770802.

Soellner L, Begemann M, Mackay DJ, Grønskov K, Tümer Z, Maher ER, Temple IK, Monk D, Riccio A, Linglart A, Netchine I, Eggermann T. Recent advances in imprinting disorders. *Clin Genet.* 2017;91:3–13. PubMed PMID: 27363536.

Takatani R, Minagawa M, Molinaro A, Reyes M, Kinoshita K, Takatani T, Kazukawa I, Nagatsuma M, Kashimada K, Sato K, Matsushita K, Nomura F, Shimojo N, Jüppner H. Similar frequency of paternal uniparental disomy involving chromosome 20q (patUPD20q) in Japanese and Caucasian patients affected by sporadic pseudohypoparathyroidism type 1b (sporPHP1B). *Bone.* 2015;79:15–20. PubMed PMID: 25997889.

## License

GeneReviews® chapters are owned by the University of Washington. Permission is hereby granted to reproduce, distribute, and translate copies of content materials for noncommercial research purposes only, provided that (i) credit for source (<http://www.genereviews.org/>) and copyright (© 1993-2024 University of Washington) are included with each copy; (ii) a link to the original material is provided whenever the material is published elsewhere on the Web; and (iii) reproducers, distributors, and/or translators comply with the [GeneReviews® Copyright Notice and Usage Disclaimer](#). No further modifications are allowed. For clarity, excerpts of GeneReviews chapters for use in lab reports and clinic notes are a permitted use.

For more information, see the [GeneReviews® Copyright Notice and Usage Disclaimer](#).

For questions regarding permissions or whether a specified use is allowed, contact: [admasst@uw.edu](mailto:admasst@uw.edu).