

Probe Report

Title: A small molecule inhibitor of Caspase 1

Authors: Matthew B. Boxer,^a Min Shen,^a Douglas S. Auld,^a James A. Wells,^b Craig J. Thomas^a

^a NIH Chemical Genomics Center, National Human Genome Research Institute, National Institutes of Health, 9800 Medical Center Drive, MSC 3370 Bethesda, Maryland 20850.

^b Departments of Pharmaceutical Chemistry and Cellular and Molecular Pharmacology, University of California, San Francisco, Byers Hall, 1700 4th Street, San Francisco, CA 94158, USA.

Assigned Assay Grant #: X01 MH078950-01

Screening Center Name & PI: NIH Chemical Genomics Center & Dr. Christopher P. Austin

Chemistry Center Name & PI: NIH Chemical Genomics Center & Dr. Christopher P. Austin


Assay Submitter & Institution: Dr. Jim Wells, University of California at San Francisco

PubChem Summary Bioassay Identifier (AID): **2389**

Abstract:

A nitrile-containing propionic acid moiety as an electrophile for covalent attack by the active site cysteine residue of caspase 1 was investigated. Several cyanopropanate containing small molecules were synthesized, including one based upon the optimized peptidic scaffold of the prodrug VX-765. A number of these compounds were potent inhibitors of caspase 1 ($IC_{50}s \leq 1$ nM). Examination of these small molecules versus a caspase panel demonstrated an impressive degree of selectivity for caspase 1 inhibition. The small molecular probe ML132 (CID-4462093; NCGC-00183434) is the most potent caspase 1 inhibitor reported to date. It also possesses a unique selectivity pattern relative to other reported caspase inhibitors. A number of these compounds were assessed for their hydrolytic stability and selected ADME properties.

Probe Structure & Characteristics:




ML132

CID/ML#	Target Name	IC ₅₀ /EC ₅₀ (nM) [SID, AID]	Anti-target Name(s)	IC ₅₀ /EC ₅₀ (μM) [SID, AID]	Fold Selective	Secondary Assay(s) Name: IC ₅₀ /EC ₅₀ (nM) [SID, AID]
44620939/ ML132	Caspase 1	0.023nM [87544173, 2389]	Panel of 9 Caspases	> 1μM [87544173, 2389]	> 1000-fold	Protease panel [87544173, 2389]

Recommendations for scientific use of the probe:

Caspase 1, also known as interleukin-converting enzyme or ICE, is responsible for the proteolytic activation of interleukin (IL)-1 β and IL-18.¹ IL-1 β and IL-18 are cytokines that play a major role in the immune response and within numerous autoimmune and inflammatory diseases. Caspase 1 is constitutively and inducibly expressed in immune response elements such as T cells, macrophages and neutrophils. Inhibitors of caspase 1 are sought for intervention strategies within ischemic disorders, Huntington's disease, amyotrophic lateral sclerosis (ALS), rheumatoid arthritis, osteoarthritis, inflammatory bowel disease and sepsis.



1. Introduction

Caspases are cysteine proteases with a strict specificity for cleaving peptide sequences C-terminal to aspartic acids residues. Currently, 12 caspase isozymes have been identified in humans, with numerous reported activities. Caspases are often subcategorized as either pro-apoptotic or pro-inflammatory enzymes. A prominent member of the pro-inflammatory class is caspase 1, also known as interleukin-converting enzyme or ICE, which is responsible for the proteolytic activation of interleukin (IL)-1 β and IL-18¹. IL-1 β and IL-18 are cytokines that play a major role in the immune response and within numerous autoimmune and inflammatory diseases². Caspase 1 is constitutively and inducibly expressed in immune response elements such as T cells, macrophages and neutrophils. Pro-caspase 1 is known to associate with several multi-protein complexes capable of responding to numerous external stimuli, suggesting that caspase 1 is a major regulator of the inflammation response. Targeting proteases, and specifically caspases, via small molecule therapeutics is an active area of research. Small molecule inhibitors of selected proteases have entered the clinic, and many have received approval. Inhibitors of caspase 1 are sought for intervention strategies within ischemic disorders, Huntington's disease, amyotrophic lateral sclerosis (ALS), rheumatoid arthritis, osteoarthritis, inflammatory bowel disease and sepsis.

The overall goal of this project is to screen for and/or design inhibitors of caspase 1 that are potent and selective over the 11 additional caspase isozymes. To date, at least three caspase 1 inhibitors have entered clinical evaluation, including Pralnacasan (VX-740), IDN-6556 and VX-765. All three agents are active site inhibitors that act through reversible (Pralnacasan and VX-765) or irreversible (IDN-6556) covalent modification of the catalytic cysteine residue. These agents have impressive potency versus caspase 1 (for instance, VX-740 has a published K_i of 1 nM) and good reported selectivity³. Any probe from this project must be an improvement on current art (i.e. potency better than VX-765 and equal selectivity). These agents, and the probes found herein, are covalent modifiers of their target. Often, covalent modifiers are avoided due to

promiscuity issues and potential for toxicities within *in vivo* studies. However, a large number of known drugs do indeed inhibit their target via covalent modification at the orthosteric site, and as tool compounds, covalent modifiers can be very useful if their selectivity can be shown versus related targets.

2 Materials and Methods

General Methods for Chemistry

All air or moisture sensitive reactions were performed under positive pressure of nitrogen with oven-dried glassware. Anhydrous solvents such as dichloromethane, *N,N*-dimethylformamide (DMF), acetonitrile, methanol and triethylamine were obtained by purchasing from Sigma-Aldrich. Preparative purification was performed on a Waters semi-preparative HPLC. The column used was a Phenomenex Luna C18 (5 micron, 30 x 75 mm) at a flow rate of 45 mL/min. The mobile phase consisted of acetonitrile and water (each containing 0.1% trifluoroacetic acid). A gradient of 10% to 50% acetonitrile over 8 minutes was used during the purification. Fraction collection was triggered by UV detection (220 nM). Analytical analysis was performed on an Agilent LC/MS (Agilent Technologies, Santa Clara, CA).

Method 1: A 7 minute gradient of 4% to 100% Acetonitrile (containing 0.025% trifluoroacetic acid) in water (containing 0.05% trifluoroacetic acid) was used with an 8 minute run time at a flow rate of 1 ml/min. A Phenomenex Luna C18 column (3 micron, 3 x 75 mm) was used at a temperature of 50°C.

Method 2: A 3 minute gradient of 4% to 100% Acetonitrile (containing 0.025% trifluoroacetic acid) in water (containing 0.05% trifluoroacetic acid) was used with a 4.5 minute run time at a flow rate of 1 mL/min. A Phenomenex Gemini Phenyl column (3 micron, 3 x 100 mm) was used at a temperature of 50 °C.

Purity determination was performed using an Agilent Diode Array Detector on both Method 1 and Method 2. Mass determination was performed using an Agilent 6130 mass spectrometer with electrospray ionization in the positive mode. ¹H NMR spectra were recorded on Varian 400 MHz

spectrometers. Chemical Shifts are reported in ppm with tetramethylsilane (TMS) as internal standard (0 ppm) for CDCl₃ solutions or undeuterated solvent (DMSO-h₆ at 2.49 ppm) for DMSO-d₆ solutions. All of the analogs for assay have purity greater than 95% based on both analytical methods. High resolution mass spectrometry was recorded on an Agilent 6210 Time-of-Flight LC/MS system. Confirmation of molecular formula was accomplished using electrospray ionization in the positive mode with the Agilent Masshunter software (version B.02).

2.1 Assays

The biochemical assay was configured using purified caspase 1 at a high enzyme concentration to promote formation of the homodimers, and thus facilitate the identification of allosteric inhibitors. The overall goal was to identify reversible inhibitors with better drug-like properties than the currently available set of aspartyl-containing peptidomimetics that covalently bind the active site. Caspase 1 was assayed using the profluorescent substrate Ac-WEHD-AFC. After initiation of the assay with substrate, the plates were rapidly read using an automated robotic system (Kalypsys, Inc.) to maintain consistent timing. A kinetic mode of detection was used where the initial rate was collected (estimated final product formation was ~10%). Compounds were screened as a concentration-titration series that ranged from 57 μM to 3.7 nM. Below is the protocol used for caspase 1.

Caspase 1 was prepared in buffer (50mM HEPES pH 7.5, 50mM KCl, 200mM NaCl, 10mM DTT, 0.1% CHAPS) at a concentration of 66.6nM, and 3 μl was dispensed to all wells using black solid Kalypsys 1536-well plates. 20nl of DMSO containing compounds was added using a Kalypsys pin-tool to columns 5-48. Then, 20nl of DMSO solution from a control plate was added to columns 1-4. Controls were: Column 1, 16 point titration with each concentration in duplicate (1:1 dilutions in DMSO; final starting concentration was 57 μM) of the caspase 1 inhibitor Ac-WEHD-CHO (Alexis Biochemicals); Column 2, a 16 point titration with each concentration in duplicate of the free AFC fluorophore prepared in DMSO (Alexis Biochemicals), final starting concentration was 40 μM; Column 3 neutral (DMSO only) control; Column 4: DMSO alone, to serve as a negative control (no substrate was added). Then 1 μl of 20 μM of the substrate Ac-WEHD-AFC (Alexis Biochemicals) prepared in the same buffer was


dispensed to all wells except columns 2 and 4, and the plates were immediately transferred (<1 min) to the Viewlux. The plates were then exposed using 405 nm excitation/520 nm emission filters for 4 sec and read at 20 sec intervals for 3 min. Final enzyme concentration was 50 nM and the final substrate concentration was 5 μ M.

Concentration-response curves were fitted to the data calculated from slope of the linear regression of fluorescent intensity versus time (the rate). The concentration-effect curves were then classified based on curve quality (r^2), response magnitude and degree of measured activity. Active compounds showed concentration-dependent decreases in the measured rate. Inconclusive compounds had appreciable concentration-dependent effects on both the measured rate and the interpolated basal fluorescence intensity at the start of the reaction, but where the basal fluorescent intensity was marginal (e.g. < 10-fold). Inconclusive compounds also encompass highly fluorescent compounds (> 100-fold increases in fluorescent intensities values relative to controls) where artificial concentration-dependent decreases in the rate occurred at high compound concentrations due to fluorescent interferences.

The assay was well behaved; the signal-to-background ratio was on average 13 and the average Z' screening factor associated with each plate was 0.8, indicating a robust performance of the screen. A total of 20 actives were found and submitted. None of these agents had the requirements to satisfy the probe definition. Additional assays were performed by commercial vendor (www.reactionbiology.com); these assays rely upon a peptide cleavage assay of a caged fluorophore (Z-LEHD)₂-R110 tetrapeptide.


2.2 Probe Chemical Characterization

Scheme 1



Conditions and reagents: (a) EDC, DMAP, CH₂Cl₂, 6 h (59%); (b) TFAA, DIPEA, CH₂Cl₂, 0 °C, 30 min. (90%); (c) thionyl chloride (excess), EtOH, 0 °C (95%); (d) TFAA, DIPEA, CH₂Cl₂, 0 °C, 30 min. (86%); (e) NMM, DME, then NH₄OH (73%); (f) TMSN₃, Bu₂SnO (0.6 equiv.), toluene, μW, 100 °C, 1h (77%); (g) TFAA, DIPEA, CH₂Cl₂, 0 °C, 30 min. (80%).

Scheme 2



Conditions and reagents: (a) EDC, HOBt, DMF, rt, 8 h; (b) DBU, CH₂Cl₂, rt (71% over 2 steps); (c) HATU, DIPEA, DMF, rt, 2 h; (d) TFA, CH₂Cl₂ (1:1), rt, 4 h (85% over 2 steps); (e) DBU, DMF, 5 min. then **15**, HATU, DIPEA, DMF, 0 °C, 2h; (f) TBAF, THF, 0 °C (72% over 2 steps); (g) DBU, DMF, 5 min. then **15**, HATU, DIPEA, DMF, 0 °C, 2h (91%); (h) DBU, DMF, 5 min. then **15**, HATU, DIPEA, DMF, 0 °C, 2h (82%).

NCGC00183434/CID:44620939/ML132: ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.68 (d, *J*=7.4 Hz, 1 H), 7.79 (d, *J*=1.9 Hz, 1 H), 7.64 (d, *J*=9.0 Hz, 1 H), 7.56 (dd, *J*=8.6, 1.9 Hz, 1 H), 6.73 (d, *J*=8.6 Hz, 1 H), 5.95 (b, 2H), 4.83 (q, *J*=7.4 Hz, 1H), 4.64 (d, *J*=9.0 Hz, 1 H), 4.20 - 4.30 (dd, *J*=8.2, 5.6 Hz, 1 H), 3.69 - 3.80 (m, 1 H), 3.60 (m, 1 H), 2.79 (dd, *J*=6.7, 3.1 Hz, 2 H), 1.99 - 2.11 (m, 1 H), 1.63 - 1.96 (m, 4 H), 0.99 (s, 9 H) ¹³C NMR (100 MHz, DMSO-*d*₆): 172.2, 170.6, 169.9, 165.8, 148.0, 129.4, 128.2, 122.1, 119.1, 116.4, 114.4, 59.7, 57.8, 48.2, 37.4, 36.6, 35.3, 29.5, 27.0, 25.1. LC/MS: Method 1, retention time: 4.595 min; Method 2, retention time: 3.591 min; HRMS: *m/z* (M+H⁺) = 477.1784 (Calculated for C₂₂H₂₈N₅O₅Cl = 477.1779). [α]_D²² = -65 (*c* 1.0, MeOH). Solubility (PBS, pH 7.4, 23°C) ≥100 μM. Stability profile over 100 hrs (water, pH 7, pH 2 and pH 8, 23°C) is shown below.




Figure 1. Aqueous stability of prodrugs VX-765 (**1**)(♦) and NCGC00185682 (**3**)(X) and drugs VRT-043198 (**2b**)(▲), NCGC00183434/CID:44620939/ML132 (**4**)(■) and NCGC00183681 (**16**)(●) at neutral (pH 7 - black), acidic (pH 2 - red), and basic (pH 8 - blue) conditions. NCGC00183434/CID:44620939 shows pH dependent stability (100% at pH=8, 40% at pH=7, and 30% at pH=2).

Probe	NCGC00183434	MLS003178557
Analog	NCGC00241069	MLS003178558
Analog	NCGC00241070	MLS003178559
Analog	NCGC00241071	MLS003178560
Analog	NCGC00185682	MLS003178561
Analog	NCGC00241073	MLS003178562

2.3 Probe Preparation

Appropriately substituted ethyl-3-cyanopropanoate derivatives are not commercially available, and therefore required synthetic elaboration. As we desired to explore both an active and prodrug form of our conceived molecule, we examined alternative protecting group strategies for the acid side chain (Scheme 1). Commercially available Fmoc protected D-isoasparagine (**5**) offered a convenient entry point to both required building blocks. Treatment of **5** with 2-(trimethylsilyl)ethanol, EDC and DMAP in methylene chloride provided the TMSE protected **6** in good yield. Conversion of **6** to nitrile **7** was accomplished by treatment with trifluoroacetic anhydride and Hunig's base. A similar sequence was used to produce the ethyl ester **9**. In addition to the ester prodrug and the active cyanopropionic acid, it was of interest to explore carboxylic acid mimetics. As such, we undertook the synthesis of a tetrazole version of the key ethyl-3-cyanopropanoate moiety. Here, we utilized the previously reported Fmoc protected (S)-2-amino-3-cyanopropanoic acid (**10**). Conversion to amide **11** was required prior to formation of

the tetrazole **12**. The amide was formed via the mixed anhydride, followed by treatment with ammonium hydroxide. Tetrazole formation was accomplished via microwave irradiation of the nitrile **11** and TMS-azide in the presence of dibutylstannane. Dehydration to nitrile **13** was accomplished in a manner analogous to **7** and **9**. With appropriately substituted/protected cyanopropanoate building blocks, we next turned our attention to the trimer core of VX-765 (Scheme 2). Both Fmoc protected L-*tert*-leucine and *tert*-butyl-L-prolinate are commercially available and were easily coupled via treatment with EDC and HOBt. Fmoc removal was effected by treatment with DBU resulting in the protected dimer **14**. Coupling of **14** with 4-amino-3-chlorobenzoic acid was accomplished using HATU and Hunig's base in DMF. TFA mediated removal of the *tert*-butyl group yielded the carboxylic acid **15**. A single pot deprotection-coupling sequence was used to generate the desired final products. Treatment of **7**, **9** and **13** with DBU in DMF effected deprotection to the free amines which were added sequentially to **15**, Hunig's base and finally HATU to yield the coupled products. The generation of **4** (NCGC00183434/CID:44620939/ML132) further required TBAF mediated removal of the TMS group.

3 Results

Please see subsection for a detailed description of the results.

3.1 Summary of Screening Results

Not applicable (these agents were rationally designed following our inability to discover new agents within the qHTS Assay for Allosteric/Competitive Inhibitors of Caspase-1; AID 900).

Identification of lead

The design of small molecule inhibitors of cysteine proteases relies heavily on covalent modification of the active site cysteine through reaction with the highly nucleophilic thiolate. Electrophilic ‘warhead’ moieties suitable for this modification include the aforementioned aldehyde, Michael acceptors (for instance, vinyl sulfones), α -halo ketones, epoxides and nitriles. In particular, nitrile-based cysteine protease inhibitors have found utility versus cathepsin K⁴, TbCatB⁵ and cruzain⁶. We endeavored to explore the promise of cyanopropanoates as caspase inhibitors (Figure 1B). To explore the potential of this functional moiety, we took advantage of the peptidic scaffold of VX-765. Further, we incorporated the ethyl-3- cyanopropanoate to mimic the prodrug qualities associated with VX-765 (Figure 1). VX-765 (**1**) is a prodrug that requires esterase cleavage of the 5-ethoxydihydrofuran-2(3H)-one moiety to yield the aldehyde functionality of the drug VRT-043198 (**2b**) (CID:11443029), which acts as a potent electrophile for attack by the active site cysteine thiol (Figure 1A). The remainder of the VX-765 (**1**) molecule establishes key binding contacts with caspase 1 that enhance the potency of the interaction and confer a modest degree of selectivity.




Figure 2. A. The structure of VX-765 (1) and schematic representation of esterase cleavage of the 5-ethoxydihydrofuran-2(3H)-one moiety to yield the active drug VRT-043198 (2b). **B.** The structure of NCGC00185682 (3) and putative esterase cleavage of the ethyl-3-cyanopropanoate moiety to yield active agent NCGC00183434/CID:44620939 (4).

3.2 Dose Response Curves for Probe




Figure 3. Structure of NCGC00183434/CID:44620939/ML132 and dose response curve for data acquired from the fluorescent enzyme assay using YVAD-CHO as a control and YVAD-AMC as the substrate.

3.3 Scaffold/Moiety Chemical Liabilities






NCGC00183434/CID:44620939/ML132 contains nitrile and acid moieties. The nitrile group is intended to be a covalent modifier of the target caspase 1. Often, covalent modifiers are avoided due to promiscuity issues and potential for toxicities within *in vivo* studies. However, a large number of known drugs do indeed inhibit their target via covalent modification at the orthosteric

site, and as tool compounds, covalent modifiers can be very useful if their selectivity can be shown versus related targets. We have profiled these agents for selectivity versus within a commercial panel of caspases offered by Reaction Biology Corporation. This data confirmed the potent inhibitory capacity of **2b** versus caspase 1 ($IC_{50} = 0.204\text{nM}$); however, the IC_{50} values found versus caspase 4 ($IC_{50} = 14.5\text{nM}$) and caspase 8 ($IC_{50} = 3. \text{nM}$) differed slightly from the reported K_i values. The results against caspase 6 ($IC_{50} \geq 10,000\text{nM}$) and caspase 9 ($IC_{50} = 5.07\text{nM}$) were significantly different from those reported by Randle and coworkers⁷. The Reaction Biology Corporation panel also included caspase 5 ($IC_{50} = 10.6\text{nM}$), caspase 10 ($IC_{50} = 66.5\text{nM}$) and caspase 14 ($IC_{50} = 58.5\text{nM}$), and the data presented here represents the first disclosure of the IC_{50} values for **2b** versus these targets. The results for **4** demonstrated an impressive potency against caspase 1 ($IC_{50} = 0.023\text{nM}$) and a similar selectivity profile as **2b**. The only prominent divergence between the selectivity profiles of **4** and **2b** was a sharp drop in the ability to inhibit caspase 14 ($IC_{50} = 801\text{nM}$ and $IC_{50} = 58.5\text{nM}$, respectively). The caspase 1 inhibition data generated in this panel for **3** and **16** was similar to the data generated in our caspase 1 assay, with reported IC_{50} values of 43.4 nM and 2.58 nM , respectively. A particularly interesting aspect of these molecules was the high selectivity for caspase 1. NCGC00183681 (**16**) registered an IC_{50} value of 91.5nM versus caspase 9. All other activities were above the $1\mu\text{M}$ threshold. In addition to the primary molecules of this study, we were interested in establishing cyanopropanoates as general caspase directing ‘warheads’ for future utility in the search for other potent and selective small molecule inhibitors of caspases. As such, we included the general nitrile-Asp directing group into the common peptide caspase inhibitors YVAD. The resulting agent, YVAD-CN (**20**), was profiled, and the results clearly demonstrate that cyanopropanoates represent a general moiety for reversible, covalent modification of caspases. Based upon this data, we conclude that these agents do not represent promiscuous covalent protein modifiers.

The acid moiety can cause issues with membrane permeability (though not always). In order to combat this, we have provided a pro-drug version of NCGC00183434/CID:44620939.

3.4 SAR Tables

Table 1. Investigation of SAR around aryl ring

Entry	Ar	Internal ID ^a	CID	SID	IC ₅₀ (nM) ^b
1		NCGC00183434	44620939	87544173	0.47
2		NCGC00241070	46916205	99380812	0.424
3		NCGC00241069	46916204	99380811	2.09
4		NCGC00241073	46916207	99380814	0.44
5		NCGC00241071	46916206	99380813	5.36

^aAll compounds synthesized at NCGC

^bIC₅₀ values were determined at NCGC utilizing a fluorescent enzyme assay using the appropriate AFC-labeled peptide.

Table 2. IC₅₀ values for selected compounds versus caspase panel.

Compound	Caspase 1 (nM)	Caspase 3 (nM)	Caspase 4 (nM)	Caspase 5 (nM)	Caspase 6 (nM)	Caspase 7 (nM)	Caspase 8 (nM)	Caspase 9 (nM)	Caspase 10 (nM)	Caspase 14 (nM)
VRT-043198 (2b) (Drug)	0.204	> 10000	14.5	10.6	> 10000	> 10000	3.3	5.07	66.5	58.5
3 (Nitrile ester)	43.4	> 10000	> 10000	1570	> 10000	> 10000	> 10000	1610	> 10000	> 10000
4 (Nitrile acid)	0.023	> 10000	13.8	3.60	> 10000	> 10000	25.2	2.17	89.7	801
16 (Nitrile tetrazole)	2.58	> 10000	1380	1300	> 10000	> 10000	> 10000	91.5	> 10000	> 10000
20 YVAD-CN	2.16	> 10000	114	29.0	> 10000	> 10000	726	297	187	116
Ac-LEHD-CHO (standard)	15.0	ND	81.7	21.3	ND	ND	3.82	49.2	40.4	134
Ac-DEVD-CHO (standard)	ND	3.04	ND	ND	122	3.54	ND	ND	ND	ND

^aData was generated by Reaction Biology (<http://www.reactionbiology.com/>). Data is presented as an IC₅₀'s using a (Z-LEHD)2-R110 tetrapeptide substrate for caspase 1, 4, 5, 8, 9, 10, 14 and a (Z-DEVD)2-R110 tetrapeptide substrate for caspase 3, 6 and 7. Data represents the results from three separate experiments.

3.5 Cellular Activity

Cellular studies have not yet been run with these agents.

3.6 Profiling Assays

We have profiled and selected analogs for ADME properties for chosen compounds. As such, 1, 2b, 3, 4 and 16 were submitted to Cyprotex for a profile of bi-directional Caco-2 permeability, plasma protein binding (both human and rat) and microsomal stability (both human and rat) studies (Table 3). All agents possessed relatively low A to B permeability; however, the pro-drug **1** and the ester **3** had moderately better levels. The high B to A levels reported for **1** and **3** strongly suggested an active transport mechanism, and a control experiment with verapamil confirmed that these agents are substrates for Pgp efflux. Unsurprisingly, the free acids **2b** and **4** and the tetrazole **16** had significantly higher free fractions in both human and rat protein binding assays relative to the more hydrophobic prodrug **1** and ethyl ester **3**. The clearance rates (Cl_{int}) and t_{1/2} for **2b**, **3**, **4** and **16** were all moderate. The ester **3** was noted to possess a slight degree of degradation in liver microsomes without NADPH as a cofactor, suggesting a non-enzymatic related degradation mechanism. The pro-drug **1** possessed minimal ability to be metabolized by liver microsomes and a t_{1/2} of >9400 minutes. It is unknown how this extended stability affects this agent's toxicity profile.

Table 3. *In vitro* ADME properties^a for selected compounds.

Compound	Caco (A2B) ^b P _{app} (x10 ⁻⁶ cms ⁻¹)	Caco (B2A) ^b P _{app} (x10 ⁻⁶ cms ⁻¹)	Protein Binding ^c fraction unbound	Microsomal Stability ^d CL _{int} (μL/min/mg protein)	Microsomal Stability ^d t _{1/2} (min)
VX-765 (2a) (Prodrug)	0.797	32.7	0.006	0.147	9430
VRT-043198 (2b) (Drug)	ND	0.173	0.420	6.72	206
3 (Nitrile ester)	0.445	9.59	0.071	27.4	50.7
4 (Nitrile acid)	0.144	0.060	0.431	10.3	134
18 (Nitrile tetrazole)	0.130	0.193	0.243	9.38	148

^aData was generated by Cyprotex (<http://www.cyprotex.com/home/>). ^bCaco-2 permeability assay over 3 separate experiments with 2 separate internal control groups. Agents were also profiled in the presence of the known Pgp substrate verapamil and data from these experiments highly suggested that **2a** and **3** were substrates for efflux by Pgp. ^cPlasma protein binding assays were performed using an equilibrium dialysis method in 100% plasma (profiles versus human and rat plasma were obtained: see the supporting information section for rat plasma binding data). ^dMicrosomal stability was profiled alongside internal control compounds, minus NADPH and minus compound (profiles versus human and rat plasma were obtained: see the supporting information section for rat plasma binding data).

4 Discussion

A few areas of SAR were explored around NCGC00183434/CID:44620939/ML132. The first investigation was to look at replacement of the 3-chloro-4-anilino-benzene ring with more hydrophobic groups. Replacement of the *p*-aniline with a *p*-methoxy or switching to the 3,5-dichlorobenzene resulted in no significant change in activity (compare entries 1, 2 and 4 in Table 1). *O*-substitution resulted in a 5 fold loss in activity (entry 3), while the *p*-trifluoromethylbenzene analog saw a 10-fold loss in potency. Investigation around the warhead was undertaken and an ethyl ester prodrug was made *vide supra* as well as a tetrazole version, which is an established carboxylic acid mimetic. As can be seen in Table 2, the tetrazole analog had lower activity, but interestingly was slightly more selective than the acid version. As expected, the ester prodrug version (NCGC00185682) did have slightly better caco-2 permeability *vide supra* (Table 3).

4.1 Comparison to existing art and how the new probe is an improvement

NCGC00183434/CID:44620939/ML132 is the most potent caspase 1 inhibitor reported to date. It also possesses a unique selectivity pattern relative to other reported caspase inhibitors.

4.2 Mechanism of Action Studies

We examined the binding mechanism of these agents through molecular modeling. Several crystal structures of caspase 1 exist, including structures with reversible and non-reversible inhibitors (PDB codes: 1BMQ, 1IBC, 1ICE, 1RWK, 1RWM, 1RWN, 1RWO, 1RWP, 1RWV, 1RWW, 1RWX, 1SC1, 1SC3, 1SC4, 2FQQ, 2H48, 2HBQ, 2HBR, 2HBY, 2HBZ, 2FQR, 2FQS, 2FQU, 2FQZ)⁸⁻¹¹. We identified 2HBQ as the best template for **4** (2HBQ is a co-crystal of caspase 1 and *Z-VAD-FMK*). We applied the presumption of a covalent reversible mechanism of inhibition when building a model for binding of **4**. The nitrile carbon was therefore held at a proximal distance (2.6 Å) from the catalytic cysteine residue (C285) by constraint docking, and flexibility was granted to the remainder of the small molecule to achieve an optimal binding pose using FRED¹². The results are shown in figure 4 and demonstrate complementarity between the peptidic fragment of **4** and the peptide binding domain of caspase 1. Key interactions were noted

for the acid moiety and arginine residues 341 and 179 in similar fashion to other Asp containing small molecule caspase 1 inhibitors. While direct interrogation of a covalent interaction between the nitrile and C285 was not pursued in our model, this representation does illustrate the open binding cavity that accommodates the tetrahedral intermediate that forms as a result of covalent binding with aldehyde based inhibitors (a mimetic of the hemithiolacetal intermediate associated with transition state 1 (TS1) during proteolysis). In contrast, covalent interactions between a thiol and a nitrile form a thioimide intermediate that mimics transition state 2 (TS2) of an enzymatic proteolytic event between a cysteine proteases and a substrate. Ménard and coworkers examined aldehyde and nitrile inhibitors of papain and found that the thioimide intermediate engages the oxyanion hole interaction in a manner that more closely mimics the natural process of hydrolysis during proteolysis¹³. This may have consequences for both the binding affinity of nitrile-based cysteine proteases inhibitors and their ultimate resolution through hydrolysis of the thioimide intermediate.




Figure 4. Molecular model (ribbon and space filling) of NCGC00183434/CID:44620939/ML132 (4) bound to caspase 1.

4.3 Planned Future Studies

Recently, the Petsko and Ringe labs have identified the human enzyme caspase-1 as the protease that cleaves α -synuclein *in vivo*, producing a C-terminal truncated fragment that nucleates the aggregation of α S. Synucleinopathies are a class of neurologic disorders that together afflict millions of Americans. They are characterized by neurocytoplasmic inclusions containing primarily α -synuclein, an abundant neuronal protein whose cellular functions are poorly understood. The most prevalent synucleinopathy is Parkinson's disease, a devastating

neurodegenerative condition that kills dopaminergic neurons, leading to tremor, slowness of movement, and eventual paralysis and death. We plan to continue our optimization of the existing probe compound in a manner that will allow it to penetrate the blood-brain barrier, thus allowing the Wells and Petsko labs to examine the role of caspase 1 in synucleinopathies, including Parkinson's disease.

Probe properties

Properties Computed from Structure

Calculated Property	Probe Identity
	CID_44620939 (MLS003178557)
Molecular Weight [g/mol]	477.94122
Molecular Formula	C22H28CIN5O5
XLogP3-AA	1.6
H-Bond Donor	4
H-Bond Acceptor	7
Rotatable Bond Count	8
Tautomer Count	10
Exact Mass	477.177897
MonoIsotopic Mass	477.177897
Topological Polar Surface Area	166
Heavy Atom Count	33
Formal Charge	0
Isotope Atom Count	0
Defined Atom StereoCenter Count	3
Undefined Atom StereoCenter Count	0
Defined Bond StereoCenter Count	0
Undefined Bond StereoCenter Count	0
Covalently-Bonded Unit Count	1
Complexity	820

5 References

- (1) F. Martinon, J. Tschopp, *Cell* **2004**, *117*, 561-574.
- (2) M. Braddock, A. Quinn, *Nat. Rev. Drug Discovery* **2004**, *3*, 1-10.
- (3) S. D. Linton, *Curr. Top. Med. Chem.* **2005**, *5*, 1697-1717.
- (4) J.-P. Falguyret, R. M. Oballa, O. Okamoto, G. Wesolowski, Y. Aubin, R. M. Ryzewski, P. Prasit, D. Riendeau, S. B. Rodan, M. D. Percival, *J. Med. Chem.* **2001**, *44*, 94-104.
- (5) J. P. Mallari, A. A. Shelat, T. Obrien, C. R. Caffrey, A. Kosinski, M. Connelly, M. Harbut, D. Greenbaum, J. H. McKerrow, R. K. Guy, *J. Med. Chem.* **2008**, *51*, 545-552.
- (6) B. T. Mott, R. Ferreira, A. Simeonov, A. Jadhav, K.-H. Ang, W. Leister, M. Shen, J. T. Silveira, J. H. McKerrow, J. Inglese, C. P. Austin, C. J. Thomas, B. K. Shoichet, D. J. Maloney, *J. Med. Chem.* **2010**, *53*, 52-60.
- (7) W. Wannamaker, R. Davies, M. Namchuk, J. Pollard, P. Ford, G. Ku, C. Decker, P. Charifson, P. Weber, U. A. Germann, K. Kuida, J. C. R. Randle, *J. Pharmacol. Exp. Ther.* **2007**, *321*, 509-516.
- (8) K. P. Wilson, J.-A. F. Black, J. A. Thomason, E. E. Kim, J. P. Griffith, M. A. Navia, M. A. Murcko, S. P. Chambers, R. A. Aldape, S. A. Raybuck, D. J. Libingston, *Nature*, **1994**, *370*, 270-275.
- (9) Y. Okamoto, H. Anan, E. Nakai, K. Morihira, Y. Yonetoku, H. Kurihara, H. Sakashita, Y. Terai, M. Takeuchi, T. Shibnuma, Y. Isomura, *Chem. Pharm. Bull.* **1999**, *47*, 11-21.
- (10) M. J. Romanowski, J. M. Scheer, T. O'Brien, R. S. McDowell, *Structure* **2004**, *12*, 1361-1371.
- (11) J. M. Scheer, M. J. Romanowski, J. A. Wells, *Proc. Nat. Acad. Sci. U. S. A.* **2006**, *103*, 7595-7600.
- (12) OpenEye Scientific Software, Inc.: Santa Fe, NM. <http://www.eyesopen.com/>
- (13) E. Dufour, A. C. Storer, R. Ménard, *Biochemistry* **1995**, *34*, 9136-9143.