

9. References

1. Global strategy for women's, children's and adolescents' health (2016–2030). New York (NY): Every Woman Every Child; 2015.
2. Strategies toward ending preventable maternal mortality. Geneva: World Health Organization; 2015.
3. Every Newborn: an action plan to end preventable deaths. Geneva: World Health Organization; 2014.
4. Comprehensive implementation plan on maternal, infant and young child nutrition. Geneva: World Health Organization; 2014.
5. World Health Organization (WHO), United Nations Children's Fund, World Bank Group. Nurturing care for early childhood development: a framework for helping children survive and thrive to transform health and human potential. Geneva: WHO; 2018.
6. WHO technical consultation on postpartum and postnatal care. Geneva: World Health Organization; 2010.
7. Gon G, Leite A, Calvert C, Woodd S, Graham WJ, Filippi V. The frequency of maternal morbidity: a systematic review of systematic reviews. *Int J Gynaecol Obstet.* 2018;141(Suppl.1):20–38. doi:10.1002/ijgo.12468.
8. Kassebaum NJ, Bertozzi-Villa A, Coggeshall MS, Shackelford KA, Steiner C, Heuton KR, et al. Global, regional, and national levels and causes of maternal mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2014;384(9947):980–1004. doi:10.1016/S0140-6736(14)60696-6.
9. Hug L, Alexander M, You D, Alkema L. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: a systematic analysis. *Lancet Glob Health.* 2019;7(6):e710–20. doi:10.1016/S2214-109X(19)30163-9.
10. United Nations Children's Fund (UNICEF), World Health Organization, World Bank Group and United Nations Inter-agency Group for Child Mortality Estimation. Levels and trends in child mortality: report 2020. New York (NY): UNICEF; 2020.
11. Campbell OM, Cegolon L, Macleod D, Benova L. Length of stay after childbirth in 92 countries and associated factors in 30 low- and middle-income countries: compilation of reported data and a cross-sectional analysis from nationally representative surveys. *PLoS Med.* 2016;13(3):e1001972. doi:10.1371/journal.pmed.1001972.
12. Requejo J, Diaz T, Park L, Chou D, Choudhury A, Guthold R, et al. Assessing coverage of interventions for reproductive, maternal, newborn, child, and adolescent health and nutrition. *BMJ.* 2020;368:l6915. doi:10.1136/bmj.l6915.
13. Ending preventable newborn deaths and stillbirths by 2030. Moving faster towards high-quality universal health coverage in 2020–2025. New York (NY) and Geneva: United Nations Children's Fund, World Health Organization; 2020.
14. United Nations Sustainable Development Goals [website]. Sustainable Development Goals (<https://www.un.org/sustainabledevelopment/sustainable-development-goals/>, accessed 20 January 2022).
15. WHO recommendations on postnatal care of the mother and newborn. Geneva: World Health Organization; 2014.
16. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva: World Health Organization; 2016.
17. WHO recommendations on intrapartum care for a positive childbirth experience. Geneva: World Health Organization; 2018.

18. Postnatal care NICE guideline NG194. National Institute for Health and Care Excellence; 2021.
19. WHO handbook for guideline development. Geneva: World Health Organization; 2014.
20. Standards for improving quality of maternal and newborn care in health facilities. Geneva: World Health Organization; 2016.
21. Finlayson K, Crossland N, Bonet M, Downe S. What matters to women in the postnatal period: a meta-synthesis of qualitative studies. *PLoS One*. 2020;15(4):e0231415. doi:10.1371/journal.pone.0231415.
22. Harvey C, Movsisyan A, Portela A, Smith H. Stakeholder perspectives on discharge preparation and discharge readiness after birth: a qualitative evidence synthesis (in preparation).
23. Alonso-Coello P, Oxman AD, Moberg J, Brignardello-Petersen R, Akl EA, Davoli M, et al.; GRADE Working Group. GRADE Evidence to Decision (EtD) frameworks: a systematic and transparent approach to making well informed healthcare choices. 2: Clinical practice guidelines. *BMJ*. 2016;353:i2089. doi:10.1136/bmj.i2089.
24. State of inequality: reproductive, maternal, newborn and child health. Geneva: World Health Organization; 2015.
25. GRADE Working Group [website] (<http://gradeworkinggroup.org/>, accessed 20 January 2022).
26. GRADE-CERQual Project Group [website] (<https://www.cerqual.org>, accessed 20 January 2022).
27. Evers S, Goossens M, de Vet H, van Tulder M, Ament A. Criteria list for assessment of methodological quality of economic evaluations: consensus on health economic criteria. *Int J Technol Assess Health Care*. 2005;21(2):240-5.
28. Sacks E, Langlois E, Ziegler D, Sauve C, Downe S, Finlayson K, et al. Factors that influence uptake of routine postnatal care: a qualitative evidence synthesis (submitted).
29. Munabi-Babigumira S, Lewin S, Glenton C, Velez M, Gonçalves-Bradley DC, Bohren MA. Factors that influence the provision of postnatal care by health workers: a qualitative evidence synthesis [protocol]. *Cochrane Database Syst Rev*. 2021;(7):CD014790.
30. Comrie-Thomson L, Gopal P, Eddy K, Baguiya A, Gerlach N, Sauvé C, et al. How do women, men and health providers perceive interventions to influence men's engagement during pregnancy, childbirth and postnatal to improve maternal and newborn health outcomes? A qualitative evidence synthesis. *Soc Sci Med*. 2021;291:114475. doi:10.1016/j.socscimed.2021.114475.
31. Prady SL, Uphoff EP, Power M, Golder S. Development and validation of a search filter to identify equity-focused studies: reducing the number needed to screen. *BMC Med Res Methodol*. 2018;18(1):106. doi:10.1186/s12874-018-0567-x.
32. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. *Cochrane handbook for systematic reviews of interventions*. Version 6.1 (updated September 2020). Cochrane; 2020.
33. Walsh D, Downe S. Appraising the quality of qualitative research. *Midwifery*. 2006;22(2):108-19. doi:10.1016/j.midw.2005.05.004.
34. OneHealth Model: intervention treatment assumptions (draft 28 September). Geneva, Glastonbury (CT): United Nations InterAgency Working Group on Costing and the Futures Institute; 2013.
35. WHO compendium of innovative health technologies for low-resource settings. Geneva: World Health Organization; 2015.
36. Langlois ÉV, Miszkurka M, Zunzunegui MV, Ghaffar A, Ziegler D, Karp I. Inequities in postnatal care in low- and middle-income countries: a systematic review and meta-analysis. *Bull World Health Organ*. 2015;93(4):259-70G. doi:10.2471/BLT.14.140996.

37. Declarations of interest. In: About WHO – ethics [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/about/ethics/declarations-of-interest>, accessed 20 January 2022).
38. Reporting the effects of an intervention in EPOC reviews. Cochrane Effective Practice and Organisation of Care; 2018.
39. Integrated management of pregnancy and childbirth: pregnancy, childbirth, postpartum and newborn care: a guide for essential practice, third edition. Geneva: World Health Organization; 2015.
40. WHO recommendations for the prevention and treatment of postpartum haemorrhage. Geneva: World Health Organization; 2012.
41. Consolidated guidelines on HIV testing services. Geneva: World Health Organization; 2019.
42. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection. Recommendations for a public health approach, second edition. Geneva: World Health Organization; 2016.
43. WHO consolidated guidelines on tuberculosis. Module 2: screening. Systematic screening for tuberculosis disease. Geneva: World Health Organization; 2021.
44. Manresa M, Pereda A, Bataller E, Terre-Rull C, Ismail KM, Webb SS. Incidence of perineal pain and dyspareunia following spontaneous vaginal birth: a systematic review and meta-analysis. *Int Urogynecol J*. 2019;30(6):853–68. doi:10.1007/s00192-019-03894-0.
45. East CE, Dorward EDF, Whale RE, Liu J. Local cooling for relieving pain from perineal trauma sustained during childbirth. *Cochrane Database Syst Rev*. 2020;(10):CD006304.
46. Deussen AR, Ashwood P, Martis R, Stewart F, Grzeskowiak LE. Relief of pain due to uterine cramping/involution after birth. *Cochrane Database Syst Rev*. 2020;(10):CD004908.
47. Milsom I, Altman D, Cartwright R, Lapitan MC, Nelson R, Sjöström S, et al., editors. *Epidemiology of urinary incontinence (UI) and other lower urinary tract symptoms (LUTS), pelvic organ prolapse (POP) and anal incontinence (AI)*. In: Abrams P, Cardozo L, Wagg A, Wein A, editors. *Incontinence: 6th International Consultation on Incontinence 2017*; Tokyo: International Continence Society; 2017.
48. Woodley SJ, Lawrenson P, Boyle R, Cody JD, Mørkved S, Kernohan A, et al. Pelvic floor muscle training for preventing and treating urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst Rev*. 2020;(5):CD007471.
49. Zakarija-Grkovic I, Stewart F. Treatments for breast engorgement during lactation. *Cochrane Database Syst Rev*. 2020;(9):CD006946.
50. Crepinsek MA, Taylor EA, Michener K, Stewart F. Interventions for preventing mastitis after childbirth. *Cochrane Database Syst Rev*. 2020;(9):CD007239.
51. Abalos E, Gyte GML, Sguassero Y. Paracetamol/acetaminophen (single administration) for perineal pain in the early postpartum period. *Cochrane Database Syst Rev*. 2021;(1):CD008407.
52. Shepherd E, Grivell RM. Aspirin (single dose) for perineal pain in the early postpartum period. *Cochrane Database Syst Rev*. 2020;(7):CD012129.
53. Wuytack F, Smith V, Cleary BJ. Oral non-steroidal anti-inflammatory drugs (single dose) for perineal pain in the early postpartum period. *Cochrane Database Syst Rev*. 2021;(1):CD011352.
54. Yildizhan R, Yildizhan B, Sahin S, Suer N. Comparison of the efficacy of diclofenac and indomethacin suppositories in treating perineal pain after episiotomy or laceration: a prospective, randomized, double-blind clinical trial. *Arch Gynecol Obstet*. 2009;280(5):735. doi:10.1007/s00404-009-1006-3.
55. Hedayati H, Parsons J, Crowther CA. Rectal analgesia for pain from perineal trauma following childbirth. *Cochrane Database Syst Rev*. 2003;(3):CD003931.

56. Hedayati H, Parsons J, Crowther CA. Topically applied anaesthetics for treating perineal pain after childbirth. *Cochrane Database Syst Rev.* 2005;(2):CD004223.
57. Medical Sciences for Health (MSH). *International medical products price guide, 2015 edition.* Medford: MSH; 2016.
58. World Health Organization Model List of Essential Medicines, 21st list. Geneva: World Health Organization; 2019.
59. Briggs GG, Freeman RK, Yaffe SJ. *Drugs in pregnancy and lactation: a reference guide to fetal and neonatal risk, eighth edition.* Philadelphia (PA): Lippincott Williams & Wilkins; 2008.
60. Moossdorff-Steinhauser HFA, Bols EMJ, Spaanderman MEA, Dirksen CD, Weemhoff M, Nieman FHM, et al. Long-term effects of motherfit group therapy in pre-(MOTHERFIT1) and post-partum women (MOTHERFIT2) with stress urinary incontinence compared to care-as-usual: study protocol of two multi-centred, randomised controlled trials. *Trials.* 2019;20(1):237. doi:10.1186/s13063-019-3331-6.
61. Salmon VE, Hay-Smith EJC, Jarvie R, Dean S, Terry R, Frawley H, et al. Implementing pelvic floor muscle training in women's childbearing years: a critical interpretive synthesis of individual, professional, and service issues. *Neurourol Urodyn.* 2020;39(2):863-70. doi:10.1002/nau.24256.
62. Hay-Smith J, Dean S, Burgio K, McClurg D, Frawley H, Dumoulin C. Pelvic-floor-muscle-training adherence "modifiers": a review of primary qualitative studies – 2011 ICS state-of-the-science seminar research paper III of IV. *Neurourol Urodyn.* 2015;34(7):622-31. doi:10.1002/nau.22771.
63. Responsive feeding infosheet [fact sheet]. Supporting close and loving relationships. London: United Nations Children's Fund UK; 2016 (<https://www.unicef.org.uk/babyfriendly/baby-friendly-resources/relationship-building-resources/responsive-feeding-infosheet/>, accessed 20 January 2022).
64. The evidence and rationale for the UNICEF UK Baby Friendly Initiative standards. London: United Nations Children's Fund UK; 2013 (<https://www.unicef.org.uk/babyfriendly/about/evidence-and-rationale-for-the-baby-friendly-standards/>, accessed 20 January 2022).
65. Guttman N, Zimmerman DR. Low-income mothers' views on breastfeeding. *Soc Sci Med.* 2000;50(10):1457-73. doi:10.1016/S0277-9536(99)00387-1.
66. Mitra AK, Khoury AJ, Hinton AW, Carothers C. Predictors of breastfeeding intention among low-income women. *Matern Child Health J.* 2004;8(2):65-70. doi:10.1023/B:MACI.0000025728.54271.27.
67. Khosravan S, Mohammadzadeh-Moghadam H, Mohammadzadeh F, Fadafen SAK, Gholami M. The effect of Hollyhock (*Althaea Officinalis* L) leaf compresses combined with warm and cold compress on breast engorgement in lactating women: a randomized clinical trial. *J Evid Based Complementary Altern Med.* 2017;22(1):25-30. doi:10.1177/2156587215617106.
68. Dehghani M, Babazadeh R, Khadivzadeh T, Pourhoseini SA, Esmaeili H. Effect of breast oketani-massage on the severity of breast engorgement. *Iran J Obstet Gynecol Infertil.* 2017;20(5):30-8. doi:10.22038/ijogi.2017.9078.
69. Lang DL, Zhao F-L, Robertson J. Prevention of postpartum haemorrhage: cost consequences analysis of misoprostol in low-resource settings. *BMC Pregnancy Childbirth.* 2015;15:305. doi:10.1186/s12884-015-0749-z.
70. Gallos I, Williams H, Price M, Pickering K, Merriel A, Tobias A, et al. Uterotonic drugs to prevent postpartum haemorrhage: a network meta-analysis. *Health Technol Assess.* 2019;23:9. doi:10.3310/hta23090.
71. Bhagat S, Agarwal M, Roy V. Serratiopeptidase: a systematic review of the existing evidence. *Int J Surg.* 2013;11(3):209-17. doi:10.1016/j.ijisu.2013.01.010.

72. Wilson E, Woodd SL, Benova L. Incidence of and risk factors for lactational mastitis: a systematic review. *J Hum Lact.* 2020;36(4):673-86. doi:10.1177/0890334420907898.
73. Higgins PD, Johanson JF. Epidemiology of constipation in North America: a systematic review. *Am J Gastroenterol.* 2004;99(4):750-9. doi:10.1111/j.1572-0241.2004.04114.x.
74. Turawa EB, Musekiwa A, Rohwer AC. Interventions for preventing postpartum constipation. *Cochrane Database Syst Rev.* 2020;(8):CD011625.
75. Guideline: protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services. Geneva: World Health Organization; 2017.
76. Guideline: counselling of women to improve breastfeeding practices. Geneva: World Health Organization; 2018.
77. Jahanfar S, Ng CJ, Teng CL. Antibiotics for mastitis in breastfeeding women. *Cochrane Database Syst Rev.* 2013;(2):CD005458.
78. Healthy diet [fact sheet]. Geneva: World Health Organization; 2020 (<https://www.who.int/en/news-room/fact-sheets/detail/healthy-diet>, accessed 20 January 2022).
79. WHO recommendations for augmentation of labour. Geneva: World Health Organization; 2014.
80. Downe S, Finlayson K, Tunçalp Ö, Gülmezoglu AM. Provision and uptake of routine antenatal services: a qualitative evidence synthesis. *Cochrane Database Syst Rev.* 2019;(6):CD012392.
81. Breastfeeding and maternal medication. Recommendations for drugs in the eleventh WHO model list of essential drugs. Geneva: World Health Organization; 2002.
82. WHO recommendations for prevention and treatment of maternal peripartum infections. Geneva: World Health Organization; 2015.
83. Preventive chemotherapy to control soil-transmitted helminth infections in at-risk population groups. Geneva: World Health Organization; 2017.
84. WHO guideline on control and elimination of human schistosomiasis. Geneva: World Health Organization; 2022.
85. Fisher J, Cabral de Mello M, Patel V, Rahman A, Tran T, Holton S, et al. Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bull World Health Organ.* 2012;90(2):139-149G. doi:10.2471/BLT.11.091850.
86. Muzik M, Thelen K, Rosenblum KL. Perinatal depression: detection and treatment. *Neuropsychiatry.* 2011;1(2):179-95. doi:10.2217/np.10.8.
87. Andermann A, Blancquaert I, Beauchamp S, Déry V. Revisiting Wilson and Jungner in the genomic age: a review of screening criteria over the past 40 years. *Bull World Health Organ.* 2008;86(4):317-9. doi:10.2471/blt.07.050112.
88. Waqas A, Koukab A, Meraj H, Dua T, Chowdhary N, Fatima B, et al. Screening programs for common maternal mental health disorders among perinatal women: report of the systematic review of evidence. *BMC Psychiatry.* 2022;22(1):54. doi:10.1186/s12888-022-03694-9.
89. Morrell CJ, Warner R, Slade P, Dixon S, Walters S, Paley G, et al. Psychological interventions for postnatal depression: cluster randomised trial and economic evaluation. The PoNDER trial. *Health Technol Assess.* 2009;13(30):iii-iv,xi-xiii,1-153. doi:10.3310/hta13300.
90. World Health Organization (WHO), Calouste Gulbenkian Foundation. Social determinants of mental health. Geneva: WHO; 2014.
91. Smith HJJ, Portela AG, Harvey C. Discharge preparation and readiness after birth: a scoping review of global policies, guidelines and literature. *BMC Pregnancy Childbirth* (in press).

92. Armstrong SJ, Small RE. The paradox of screening: rural women's views on screening for postnatal depression. *BMC Public Health*. 2010;10(1):744. doi:10.1186/1471-2458-10-744.
93. Buist A, Condon J, Brooks J, Speelman C, Milgrom J, Hayes B, et al. Acceptability of routine screening for perinatal depression. *J Affect Disord*. 2006;93(1-3):233-7. doi:10.1016/j.jad.2006.02.019.
94. Waqas A, Kokab A, Meraj H, Dua T, Chowdhary N, Fatima B, et al. Prevention of common mental disorders among women in the perinatal period: a critical mixed-methods review and meta-analysis. *Glob Ment Health* (in press).
95. Dukhovny D, Dennis CL, Hodnett E, Weston J, Stewart DE, Mao W, et al. Prospective economic evaluation of a peer support intervention for prevention of postpartum depression among high-risk women in Ontario, Canada. *Am J Perinatol*. 2013;30(8):631-42. doi:10.1055/s-0032-1331029.
96. Ride J, Lorgelly P, Tran T, Wynter K, Rowe H, Fisher J. Preventing postnatal maternal mental health problems using a psychoeducational intervention: the cost-effectiveness of What Were We Thinking. *BMJ Open*. 2016;6(11):e012086. doi:10.1136/bmjopen-2016-012086.
97. Delaney AM, George Dalmida S, Gaydos L. When it is not postpartum depression: understanding the postpartum mental health needs of mothers and how they are being met. *Issues Ment Health Nurs*. 2015;36(6):416-23. doi:10.3109/01612840.2014.1002645.
98. Raymond NC, Pratt RJ, Godecker A, Harrison PA, Kim H, Kuendig J, et al. Addressing perinatal depression in a group of underserved urban women: a focus group study. *BMC Pregnancy Childbirth*. 2014;14:336. doi:10.1186/1471-2393-14-336.
99. Thurtle V. First time mothers' perceptions of motherhood and PND. *Community Practitioner*. 2003;76:261-5.
100. Iron supplementation in postpartum women. Geneva: World Health Organization; 2016.
101. Daily iron supplementation in adult women and adolescent girls. Geneva: World Health Organization; 2016.
102. Intermittent iron and folic acid supplementation in menstruating women. Geneva: World Health Organization; 2011.
103. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva: World Health Organization; 2011.
104. International Nutritional Anemia Consultative Group, World Health Organization, United Nations Children's Fund. Guidelines for the use of iron supplements to prevent and treat iron deficiency anaemia. Washington (DC): ILSI Press; 1998.
105. Healthy eating during pregnancy and breastfeeding. Booklet for mothers. Copenhagen: World Health Organization; 2001.
106. Vitamin A supplementation in postpartum women. Geneva: World Health Organization; 2011.
107. WHO guidelines on physical activity and sedentary behaviour. Geneva: World Health Organization; 2020.
108. Ensuring human rights in the provision of contraceptive information and services: guidance and recommendations. Geneva: World Health Organization; 2014.
109. Medical eligibility criteria for contraceptive use. Geneva: World Health Organization; 2015.
110. Contraceptive eligibility for women at high risk of HIV. Geneva: World Health Organization; 2019.
111. WHO consolidated guideline on self-care interventions for health: sexual and reproductive health and rights. Geneva: World Health Organization; 2019.
112. Rahi J, Gilbert CE. Epidemiology and the worldwide impact of visual impairment in children. In: Lyons CJ, Lambert SR, editors. *Taylor and Hoyt's Pediatric Ophthalmology and Strabismus*, sixth edition [e-book]. London, New York, Oxford, Philadelphia, St Louis, Sydney, Toronto: Elsevier; 2017.

113. Malik ANJ, Evans JR, Gupta S, Mariotti S, Gordon I, Bowman R, et al. Universal newborn eye screening: a systematic review of the literature and review of international guidelines (submitted).
114. Lieu JEC, Kenna M, Anne S, Davidson L. Hearing loss in children: a review. *JAMA*. 2020;324(21):2195–205. doi:10.1001/jama.2020.17647.
115. Butcher E, Dezateux C, Cortina-Borja M, Knowles RL. Prevalence of permanent childhood hearing loss detected at the universal newborn hearing screen: Systematic review and meta-analysis. *PLoS One*. 2019;14(7):e0219600. doi:10.1371/journal.pone.0219600.
116. World report on hearing. Geneva: World Health Organization; 2021.
117. Bhutani VK, Zipursky A, Blencowe H, Khanna R, Sgro M, Ebbesen F, et al. Neonatal hyperbilirubinemia and Rhesus disease of the newborn: incidence and impairment estimates for 2010 at regional and global levels. *Pediatr Res*. 2013;74 Suppl 1(Suppl 1):86–100. doi:10.1038/pr.2013.208.
118. Olusanya BO, Kaplan M, Hansen TWR. Neonatal hyperbilirubinaemia: a global perspective. *Lancet Child Adolesc Health*. 2018;2(8):610–20. doi:10.1016/s2352-4642(18)30139-1.
119. Sun M, Ma A, Li F, Cheng K, Zhang M, Yang H, et al. Sensitivity and specificity of red reflex test in newborn eye screening. *J Pediatr*. 2016;179:192–6.e4. doi:10.1016/j.jpeds.2016.08.048.
120. Haargaard B, Nyström A, Rosensvärd A, Tornqvist K, Magnusson G. The Pediatric Cataract Register (PECARE): analysis of age at detection of congenital cataract. *Acta Ophthalmol*. 2015;93(1):24–6. doi:10.1111/aos.12445.
121. Magnusson G, Persson U. Screening for congenital cataracts: a cost-consequence analysis of eye examination at maternity wards in comparison to well-baby clinics. *Acta Paediatr*. 2005;94(8):1089–95. doi:10.1111/j.1651-2227.2005.tb02050.x.
122. American Academy of Pediatrics. Red reflex examination in neonates, infants, and children. *Pediatrics*. 2008;122(6):1401–4. doi:10.1542/peds.2008-2624.
123. UNICEF supply catalogue [website]. United Nations Children's Fund; 2018 (<https://supply.unicef.org>, accessed 20 January 2022).
124. Lowe J, Cleland CR, Mgaya E, Furahini G, Gilbert CE, Burton MJ, et al. The Arclight ophthalmoscope: a reliable low-cost alternative to the standard direct ophthalmoscope. *J Ophthalmol*. 2015;2015:743263. doi:10.1155/2015/743263.
125. Kong L, Fry M, Al-Samarraie M, Gilbert C, Steinkuller PG. An update on progress and the changing epidemiology of causes of childhood blindness worldwide. *JAAPOS*. 2012;16(6):501–7. doi:10.1016/j.jaapos.2012.09.004.
126. Gilbert CE, Lepvrier-Chomette N. Gender inequalities in surgery for bilateral cataract among children in low-income countries: a systematic review. *Ophthalmology*. 2016;123(6):1245–51. doi:10.1016/j.optha.2016.01.048.
127. Fabian ID, Abdallah E, Abdullahi SU, Abdulqader RA, Adamou Boubacar S, Ademola-Popoola DS, et al.; Global Retinoblastoma Study Group. Global retinoblastoma presentation and analysis by national income level. *JAMA Oncol*. 2020;6(5):685–95. doi:10.1001/jamaoncol.2019.6716.
128. Malik ANJ, Mafwiri M, Gilbert C, Kim MJ, Schellenberg J. Integrating eye health training into the primary child healthcare programme in Tanzania: a pre-training and post-training study. *BMJ Paediatr Open*. 2020;4(1):e000629. doi:10.1136/bmjpo-2019-000629.
129. Mndeme FG, Mmbaga BT, Kim MJ, Sinke L, Allen L, Mgaya E, et al. Red reflex examination in reproductive and child health clinics for early detection of paediatric cataract and ocular media disorders: cross-sectional diagnostic accuracy and feasibility studies from Kilimanjaro, Tanzania. *Eye*. 2021;35:1347–53. doi:10.1038/s41433-020-1019-5.

130. Universal Newborn Hearing Screening (UNHS) review group. Effectiveness of universal newborn hearing screening: a systematic review and meta-analysis (in preparation).
131. Nelson HD, Bougatsos C, Nygren P. Universal newborn hearing screening: systematic review to update the 2001 US Preventive Services Task Force Recommendation. *Pediatrics*. 2008;122(1):e266-76. doi:10.1542/peds.2007-1422.
132. Patel H, Feldman M, Canadian Paediatric S, Community Paediatrics C. Universal newborn hearing screening. *Paediatr Child Health*. 2011;16(5):301-5. doi:10.1093/pch/16.5.301.
133. Sharma R, Gu Y, Ching TYC, Marnane V, Parkinson B. Economic evaluations of childhood hearing loss screening programmes: a systematic review and critique. *Appl Health Econ*. 2019;17(3):331-57. doi:10.1007/s40258-018-00456-1.
134. Olusanya BO, Emokpae A, Renner JK, Wirz SL. Costs and performance of early hearing detection programmes in Lagos, Nigeria. *Trans R Soc Trop Med Hyg*. 2009;103(2):179-86. doi:10.1016/j.trstmh.2008.07.001.
135. Heidari S, Olyaei Manesh A, Rajabi F, Moradi-Joo M. Cost-effectiveness analysis of automated auditory brainstem response and otoacoustic emission in universal neonatal hearing screening. *Iran J Pediatr*. 2017;27(2):e5229. doi:10.5812/ijp.5229.
136. Gupta S, Sah S, Som T, Saksena M, Yadav CP, Sankar MJ, et al. Challenges of implementing universal newborn hearing screening at a tertiary care centre from India. *Indian J Pediatr*. 2015;82(8):688-93. doi:10.1007/s12098-015-1688-4.
137. Verkleij ML, Heijnsdijk EAM, Bussé AML, Carr G, Goedegebure A, Mackey AR, et al. Cost-effectiveness of neonatal hearing screening programs: a micro-simulation modeling analysis. *Ear Hear*. 2021;42(4):909-16. doi:10.1097/AUD.0000000000000981.
138. Coates H, Giffkins K. Newborn hearing screening. *Australian Prescriber*. 2003;26(4):82-4. doi:10.18773/austprescr.2003.062.
139. Olusanya BO, Newton VE. Global burden of childhood hearing impairment and disease control priorities for developing countries. *Lancet*. 2007;369(9569):1314-7. doi:10.1016/S0140-6736(07)60602-3.
140. Van Kerschaver E, Boudewyns AN, Declau F, Van de Heyning PH, Wuyts FL. Socio-demographic determinants of hearing impairment studied in 103 835 term babies. *Eur J Public Health*. 2012;23(1):55-60. doi:10.1093/eurpub/cks010.
141. Lantos PM, Maradiaga-Panayotti G, Barber X, Raynor E, Tucci D, Hoffman K, et al. Geographic and racial disparities in infant hearing loss. *Otolaryngol Head Neck Surg*. 2018;194599818803305. doi:10.1177/0194599818803305.
142. Frary CD, Thomsen P, Gerke O. Risk factors for non-participation in the Danish universal newborn hearing screening program: a population-based cohort study. *Int J Pediatr Otorhinolaryngol*. 2020;135:110079. doi:10.1016/j.ijporl.2020.110079.
143. Young A, Tattersall H. Parents' of deaf children evaluative accounts of the process and practice of universal newborn hearing screening. *J Deaf Stud Deaf Educ*. 2005;10(2):134-45. doi:10.1093/deafed/eni014.
144. Young A, Tattersall H. Universal newborn hearing screening and early identification of deafness: parents' responses to knowing early and their expectations of child communication development. *J Deaf Stud Deaf Educ*. 2007;12(2):209-20. doi:10.1093/deafed/enl033.
145. Arnold CL, Davis TC, Humiston SG, Bocchini JA, Jr., Bass PF, 3rd, Bocchini A, et al. Infant hearing screening: stakeholder recommendations for parent-centered communication. *Pediatrics*. 2006;117(5 Pt 2):S341-54. doi:10.1542/peds.2005-2633N.
146. Jatto ME, Ogunkeyede SA, Adeyemo AA, Adeagbo K, Saiki O. Mothers' perspectives of newborn hearing screening programme. *Ghana Med J*. 2018;52(3):158-62. doi:10.4314/gmj.v52i3.9.

147. Shulman S, Besculides M, Saltzman A, Ireys H, White KR, Forsman I. Evaluation of the universal newborn hearing screening and intervention program. *Pediatrics*. 2010;126 Suppl 1:S19-27. doi:10.1542/peds.2010-0354F.
148. Recommendations for management of common childhood conditions. Evidence for technical update of pocket book recommendations. Geneva: World Health Organization; 2012.
149. Khurshid F, Rao SPN, Sauve C, Gupta S. Universal screening for hyperbilirubinemia in term healthy newborns at discharge: a systematic review and meta-analysis (submitted).
150. Wainer S, Parmar SM, Allegro D, Rabi Y, Lyon ME. Impact of a transcutaneous bilirubinometry program on resource utilization and severe hyperbilirubinemia. *Pediatrics*. 2012;129(1):77-86. doi:10.1542/peds.2011-0599.
151. Olusanya BO, Imosemi DO, Emokpae AA. Differences between transcutaneous and serum bilirubin measurements in Black African neonates. *Pediatrics*. 2016;138(3). doi:10.1542/peds.2016-0907.
152. Taylor JA, Burgos AE, Flaherman V, Chung EK, Simpson EA, Goyal NK, et al. Discrepancies between transcutaneous and serum bilirubin measurements. *Pediatrics*. 2015;135(2):224-31. doi:10.1542/peds.2014-1919.
153. Bhutani VK, Gourley GR, Adler S, Kreamer B, Dalin C, Johnson LH. Noninvasive measurement of total serum bilirubin in a multiracial pre-discharge newborn population to assess the risk of severe hyperbilirubinemia. *Pediatrics*. 2000;106(2):E17. doi:10.1542/peds.106.2.e17.
154. McClean S, Baerg K, Smith-Fehr J, Szafron M. Cost savings with transcutaneous screening versus total serum bilirubin measurement for newborn jaundice in hospital and community settings: a cost-minimization analysis. *CMAJ Open*. 2018;6(3):E285-91. doi:10.9778/cmajo.20170158.
155. Olusanya BO, Ogunlesi TA, Slusher TM. Why is kernicterus still a major cause of death and disability in low-income and middle-income countries? *Arch Dis Child*. 2014;99(12):1117-21. doi:10.1136/archdischild-2013-305506.
156. Lipstein EA, Nabi E, Perrin JM, Luff D, Browning MF, Kuhlthau KA. Parents' decision-making in newborn screening: opinions, choices, and information needs. *Pediatrics*. 2010;126(4):696-704. doi:10.1542/peds.2010-0217.
157. Maisels MJ. Neonatal transcutaneous bilirubin measurements: an opportunity to enhance laboratory utilization and improve patient care. *Clin Chem*. 2012;58(10):1395-6. doi:10.1373/clinchem.2012.183616.
158. Xie B, da Silva O, Zaric G. Cost-effectiveness analysis of a system-based approach for managing neonatal jaundice and preventing kernicterus in Ontario. *Paediatr Child Health*. 2012;17(1):11-16. doi:10.1093/pch/17.1.11.
159. Lunze K, Bloom DE, Jamison DT, Hamer DH. The global burden of neonatal hypothermia: systematic review of a major challenge for newborn survival. *BMC Med*. 2013;11(1):24. Doi:10.1186/1741-7015-11-24.
160. Kelly PA, Classen KA, Crandall CG, Crenshaw JT, Schaefer SA, Wade DA, et al. Effect of timing of the first bath on a healthy newborn's temperature. *J Obstet Gynecol Neonatal Nurs*. 2018;47(5):608-19. doi:10.1016/j.jogn.2018.07.004.
161. Ruschel LM, Pedrini DB, da Cunha MLC. Hypothermia and the newborn's bath in the first hours of life. *Rev Gaucha Enferm*. 2018;39:e20170263. doi:10.1590/1983-1447.2018.20170263.
162. Simpson EL, Chalmers JR, Hanifin JM, Thomas KS, Cork MJ, McLean WH, et al. Emollient enhancement of the skin barrier from birth offers effective atopic dermatitis prevention. *J Allergy Clin Immunol*. 2014;134(4):818-23. doi:10.1016/j.jaci.2014.08.005.
163. Dyer JA. Newborn skin care. *Semin Perinatol*. 2013;37(1):3-7. doi:10.1053/j.semperi.2012.11.008.

164. Lawn JE, Blencowe H, Oza S, You D, Lee AC, Waiswa P, et al. Every newborn: progress, priorities, and potential beyond survival. *Lancet*. 2014;384(9938):189–205. doi:10.1016/s0140-6736(14)60496-7.
165. Imdad A, Bautista RMM, Senen KAA, Uy MEV, Mantaring Iii JB, Bhutta ZA. Umbilical cord antiseptics for preventing sepsis and death among newborns. *Cochrane Database Syst Rev*. 2013;(5):CD008635.
166. Coffey PS, Brown SC. Umbilical cord-care practices in low- and middle-income countries: a systematic review. *BMC Pregnancy Childbirth*. 2017;17(1):68. doi:10.1186/s12884-017-1250-7.
167. Krous HF, Beckwith JB, Byard RW, Rognum TO, Bajanowski T, Corey T, et al. Sudden infant death syndrome and unclassified sudden infant deaths: a definitional and diagnostic approach. *Pediatrics*. 2004;114(1):234–8. doi:10.1542/peds.114.1.234.
168. Priyadarshi M, Balachander B, Sankar MJ. Effect of sleep position in term healthy newborns on neonatal mortality and sudden infant death syndrome (SIDS): a systematic review (submitted).
169. Filiano JJ, Kinney HC. A perspective on neuropathologic findings in victims of the sudden infant death syndrome: the triple-risk model. *Biol Neonate*. 1994;65(3-4):194–7. doi:10.1159/000244052.
170. Priyadarshi M, Balachander B, Gupta S, Sankar MJ. Timing of bathing in term healthy newborns: a systematic review (submitted).
171. Callaghan-Koru JA, Nonyane BAS, Guenther T, Sitrin D, Ligowe R, Chimbalanga E, et al. Contribution of community-based newborn health promotion to reducing inequities in healthy newborn care practices and knowledge: evidence of improvement from a three-district pilot program in Malawi. *BMC Public Health*. 2013;13(1):1052. doi:10.1186/1471-2458-13-1052.
172. Nonyane BA, KC A, Callaghan-Koru JA, Guenther T, Sitrin D, Syed U, et al. Equity improvements in maternal and newborn care indicators: results from the Bardiya district of Nepal. *Health Policy Plan*. 2015;31(4):405–14. doi:10.1093/heapol/czv077.
173. Chamberlain J, McCarty S, Sorce J, Leesman B, Schmidt S, Meyrick E, et al. Impact on delayed newborn bathing on exclusive breastfeeding rates, glucose and temperature stability, and weight loss. *J Neonatal Nurs*. 2019;25:74–7. doi:10.1016/j.jnn.2018.11.001.
174. Lund C. Bathing and beyond: current bathing controversies for newborn infants. *Adv Neonatal Care*. 2016;16 Suppl 5S:S13–20. doi:10.1097/anc.0000000000000336.
175. Priyadarshi M, Balachander B, Gupta S, Sankar MJ. Emollients application in term healthy newborns: a systematic review (submitted).
176. Chalmers JR, Haines RH, Mitchell EJ, Thomas KS, Brown SJ, Ridd M, et al. Effectiveness and cost-effectiveness of daily all-over-body application of emollient during the first year of life for preventing atopic eczema in high-risk children (The BEEP trial): protocol for a randomised controlled trial. *Trials*. 2017;18(1):343. doi:10.1186/s13063-017-2031-3.
177. Chlorhexidine Umbilical Review Group. Efficacy and safety of umbilical cord cleansing with chlorhexidine in neonates – an individual participant data (IPD) meta-analysis (in preparation).
178. Information Exchange System. Alert No. 133. Chlorhexidine 7,1% digluconate (CHX) aqueous solution or gel (10ml) – reports of serious eye injury due to errors in administration. Geneva: World Health Organization; 2019.
179. Coffey PS, Metzler M, Islam Z, Koehlmoos TP. Willingness to pay for a 4% chlorhexidine (7.1% chlorhexidine digluconate) product for umbilical cord care in rural Bangladesh: a contingency valuation study. *BMC Int Health Hum Rights*. 2013;13:44. doi:10.1186/1472-698x-13-44.
180. Arifeen SE, Mullany LC, Shah R, Mannan I, Rahman SM, Talukder MR, et al. The effect of cord cleansing with chlorhexidine on neonatal mortality in rural Bangladesh: a community-based, cluster-randomised trial. *Lancet*. 2012;379(9820):1022–8. doi:10.1016/s0140-6736(11)61848-5.

181. World Health Organization Model List of Essential Medicines for Children, 7th list. Geneva: World Health Organization; 2019.
182. Willinger M, Hoffman HJ, Hartford RB. Infant sleep position and risk for sudden infant death syndrome: report of meeting held January 13 and 14, 1994, National Institutes of Health, Bethesda, MD. *Pediatrics*. 1994;93(5):814-9.
183. Cheng TL, Monteiro N, DiMeglio LA, Chien AT, Peeples ES, Raetz E, et al. Seven great achievements in pediatric research in the past 40 y. *Pediatr Res*. 2016;80(3):330-7. doi:10.1038/pr.2016.95.
184. Task Force on Sudden Infant Death Syndrome, Moon RY. SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*. 2011;128(5):e1341-67. doi:10.1542/peds.2011-2285.
185. Katz D, Shore S, Bandle B, Niermeyer S, Bol KA, Khanna A. Sudden infant death syndrome and residential altitude. *Pediatrics*. 2015;135(6):e1442-9. doi:10.1542/peds.2014-2697.
186. Spencer N, Logan S. Sudden unexpected death in infancy and socioeconomic status: a systematic review. *J Epidemiol Community Health*. 2004;58(5):366-73. doi:10.1136/jech.2003.011551.
187. Hwang SS, Smith RA, Barfield WD, Smith VC, McCormick MC, Williams MA. Supine sleep positioning in preterm and term infants after hospital discharge from 2000 to 2011. *J Perinatol*. 2016;36(9):787-93. doi:10.1038/jp.2016.80.
188. Smylie J, Fell DB, Chalmers B, Sauve R, Royle C, Allan B, et al. Socioeconomic position and factors associated with use of a nonsupine infant sleep position: findings from the Canadian Maternity Experiences Survey. *Am J Public Health*. 2014;104(3):539-47. doi:10.2105/ajph.2012.301061.
189. Hirai AH, Kortsmid K, Kaplan L, Reiney E, Warner L, Parks SE, et al. Prevalence and factors associated with safe infant sleep practices. *Pediatrics*. 2019;144(5). doi:10.1542/peds.2019-1286.
190. Zachry AH, Kitzmann KM. Disparities in sleep position awareness and compliance. *South Med J*. 2010;103(4):311-5. doi:10.1097/SMJ.0b013e3181d3ce78.
191. Pease A, Garstang JJ, Ellis C, Watson D, Ingram J, Cabral C, et al. Decision-making for the infant sleep environment among families with children considered to be at risk of sudden unexpected death in infancy: a systematic review and qualitative metasynthesis. *BMJ Paediatr Open*. 2021;5(1):e000983. doi:10.1136/bmjpo-2020-000983.
192. World Health Organization. Hepatitis B vaccines: WHO position paper. *Wkly Epidemiol Rec*. 2017;27(92):369-92.
193. World Health Organization. Polio vaccines: WHO position paper. *Wkly Epidemiol Rec*. 2016;12(91):145-68.
194. World Health Organization. BCG vaccines: WHO position paper. *Wkly Epidemiol Rec*. 2018;8(93):73-96.
195. WHO recommendations for routine immunization - summary tables [website]. Geneva: World Health Organization; 2021 (<https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/who-recommendations-for-routine-immunization---summary-tables>, accessed 20 January 2022).
196. World Health Organization (WHO), Food and Agriculture Organization of the United Nations. Vitamin and mineral requirements in human nutrition, second edition. Geneva: WHO; 2004.
197. Haider BA, Sharma R, Bhutta ZA. Neonatal vitamin A supplementation for the prevention of mortality and morbidity in term neonates in low and middle income countries. *Cochrane Database Syst Rev*. 2017;(2):CD006980.
198. Global prevalence of vitamin A deficiency in populations at risk 1995-2005. WHO global database on vitamin A deficiency. Geneva: World Health Organization; 2009.

199. Stevens GA, Bennett JE, Hennocq Q, Lu Y, De-Regil LM, Rogers L, et al. Trends and mortality effects of vitamin A deficiency in children in 138 low-income and middle-income countries between 1991 and 2013: a pooled analysis of population-based surveys. *Lancet Glob Health*. 2015;3(9):e528–36. doi:10.1016/S2214-109X(15)00039-X.
200. Koo W, Walyat N. Vitamin D and skeletal growth and development. *Curr Osteoporos Rep*. 2013;11(3):188–93. doi:10.1007/s11914-013-0156-1.
201. Winzenberg T, Jones G. Vitamin D and bone health in childhood and adolescence. *Calcif Tissue Int*. 2013;92(2):140–50. doi:10.1007/s00223-012-9615-4.
202. Nutritional rickets: a review of disease burden, causes, diagnosis, prevention and treatment. Geneva: World Health Organization; 2019.
203. Við Streyms S, Højskov CS, Møller UK, Heickendorff L, Vestergaard P, Mosekilde L, et al. Vitamin D content in human breast milk: a 9-month follow-up study. *Am J Clin Nutr*. 2016;103(1):107–14. doi:10.3945/ajcn.115.115105.
204. Imdad A, Rehman F, Davis E, Ranjit D, Surin GSS, Attia SL, et al. Effects of neonatal nutrition interventions on neonatal mortality and child health and development outcomes: a systematic review. *Campbell Syst Rev*. 2019;17(1):e1141. doi:10.1002/cl2.1021.
205. Neonatal Vitamin A Supplementation Evidence group. Early neonatal vitamin A supplementation and infant mortality: an individual participant data meta-analysis of randomised controlled trials. *Arch Dis Child*. 2019;104(3):217–26. doi:10.1136/archdischild-2018-315242.
206. Humphrey JH, Agoestina T, Juliana A, Septiana S, Widjaja H, Cerreto MC, et al. Neonatal vitamin A supplementation: effect on development and growth at 3 years of age. *Am J Clin Nutr*. 1998;68(1):109–17. doi:10.1093/ajcn/68.1.109.
207. Klemm RD, Labrique AB, Christian P, Rashid M, Shamim AA, Katz J, et al. Newborn vitamin A supplementation reduced infant mortality in rural Bangladesh. *Pediatrics*. 2008;122(1):e242–50. doi:10.1542/peds.2007-3448.
208. Edmond KM, Newton S, Shannon C, O’Leary M, Hurt L, Thomas G, et al. Effect of early neonatal vitamin A supplementation on mortality during infancy in Ghana (Neovita): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015;385(9975):1315–23. doi:10.1016/s0140-6736(14)60880-1.
209. Masanja H, Smith ER, Muhimi A, Briegleb C, Mshamu S, Ruben J, et al. Effect of neonatal vitamin A supplementation on mortality in infants in Tanzania (Neovita): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015;385(9975):1324–32. doi:10.1016/s0140-6736(14)61731-1.
210. Mazumder S, Taneja S, Bhatia K, Yoshida S, Kaur J, Dube B, et al. Efficacy of early neonatal supplementation with vitamin A to reduce mortality in infancy in Haryana, India (Neovita): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015;385(9975):1333–42. doi:10.1016/s0140-6736(14)60891-6.
211. Ahmad SM, Raqib R, Huda MN, Alam MJ, Monirujjaman M, Akhter T, et al. High-dose neonatal vitamin A supplementation transiently decreases thymic function in early infancy. *J Nutr*. 2020;150(1):176–83. doi:10.1093/jn/nxz193.
212. Klemm RDW, Pokharel RK, Bichha RP, Nargis M, Mahmud Z, Oot L, et al. Testing the feasibility of delivering vitamin A to newborns in Nepal and Bangladesh. *Sight and Life*. 2011;25:30–7.
213. Codex Alimentarius Commission. Standard on infant formula and formulas for special medical purposes intended for infants. Formerly CAC/RS 72-1972. Adopted as a worldwide standard in 1981. Geneva: World Health Organization; 2007.
214. Tan ML, Abrams SA, Osborn DA. Vitamin D supplementation for term breastfed infants to prevent vitamin D deficiency and improve bone health. *Cochrane Database Syst Rev*. 2020;(12):CD013046.
215. Optimal feeding of low birth-weight infants in low- and middle-income countries. Geneva: World Health Organization; 2011.

216. Chandy DD, Kare J, Singh SN, Agarwal A, Das V, Singh U, et al. Effect of vitamin D supplementation, directly or via breast milk for term infants, on serum 25 hydroxyvitamin D and related biochemistry, and propensity to infection: a randomised placebo-controlled trial. *Br J Nutr*. 2016;116(1):52–8. doi:10.1017/s0007114516001756. (https://www.cdc.gov/nccdphp/dnpa/nutrition/pdf/Vitamin_D_Expert_Panel_Meeting.pdf, accessed 20 January 2022).
217. Floreskul V, Juma FZ, Daniel AB, Zamir I, Rawdin A, Stevenson M, et al. Cost-effectiveness of vitamin D supplementation in pregnant woman and young children in preventing rickets: a modeling study. *Front Public Health*. 2020;8(439). doi:10.3389/fpubh.2020.00439.
218. Roth DE, Abrams SA, Aloia J, Bergeron G, Bourassa MW, Brown KH, et al. Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low- and middle-income countries. *Ann N Y Acad Sci*. 2018;1430(1):44–79. doi:10.1111/nyas.13968.
219. Wagner CL, Greer FR. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics*. 2008;122(5):1142–52. doi:10.1542/peds.2008-1862.
220. Simon AE, Ahrens KA. Adherence to vitamin D intake guidelines in the United States. *Pediatrics*. 2020;145(6):e20193574. doi:10.1542/peds.2019-3574.
221. Umaretiya PJ, Oberhelman SS, Cozine EW, Maxson JA, Quigg SM, Thacher TD. Maternal preferences for vitamin D supplementation in breastfed infants. *Ann Fam Med*. 2017;15(1):68–70. doi:10.1370/afm.2016.
222. Taylor JA, Geyer LJ, Feldman KW. Use of supplemental vitamin D among infants breastfed for prolonged periods. *Pediatrics*. 2010;125(1):105–11. doi:10.1542/peds.2009-1195.
223. Heinig MJ. Vitamin D and the breastfed infant: controversies and concerns. *J Hum Lact*. 2003;19(3):247–9. doi:10.1177/0890334403255420.
224. Vitamin D expert panel meeting final report. Atlanta, GA: Centers for Disease Control; 2001.
225. Crocker B, Green TJ, Barr SI, Beckingham B, Bhagat R, Dabrowska B, et al. Very high vitamin D supplementation rates among infants aged 2 months in Vancouver and Richmond, British Columbia, Canada. *BMC Public Health*. 2011;11(1):905. doi:10.1186/1471-2458-11-905.
226. Gallo S, Jean-Philippe S, Rodd C, Weiler HA. Vitamin D supplementation of Canadian infants: practices of Montreal mothers. *Appl Physiol Nutr Metab*. 2010;35(3):303–9. doi:10.1139/h10-021.
227. Hemmingway A, Fisher D, Berkery T, Murray DM, Kiely ME. Adherence to the infant vitamin D supplementation policy in Ireland. *Eur J Nutr*. 2020;1337–45. doi:10.1007/s00394-020-02334-w.
228. Field T. Pediatric massage therapy research: a narrative review. *Children (Basel)*. 2019;6(6). doi:10.3390/children6060078.
229. Kulkarni A, Kaushik JS, Gupta P, Sharma H, Agrawal RK. Massage and touch therapy in neonates: the current evidence. *Indian Pediatr*. 2010;47:771–6. doi:10.1007/s13312-010-0114-2.
230. Diego MA, Field T. Moderate pressure massage elicits a parasympathetic nervous system response. *Int J Neurosci*. 2009;119(5):630–8. doi:10.1080/00207450802329605.
231. Schanberg S, Davalos M, Malphurs J. Massage with oil has more positive effects on normal infants. *J Prenat Perinat Psychol Health*. 1996;11(2):75.
232. Field T. Massage therapy research review. *Complement Ther Clin Pract*. 2016;24:19–31. doi:10.1016/j.ctcp.2016.04.005.
233. Priyadarshi M, Kumar V, Balachander B, Gupta S, Sankar MJ. Effect of whole-body massage on growth and neurodevelopment in term healthy newborns: a systematic review (submitted).

234. Bennett C, Underdown A, Barlow J. Massage for promoting mental and physical health in typically developing infants under the age of six months. *Cochrane Database Syst Rev.* 2013;(4):CD005038.
235. Zhang M, Wang L, Wang Y, Tang J. The influence of massage on neonatal hyperbilirubinemia: a meta-analysis of randomized controlled trials. *J Matern-Fetal Neonatal Med.* 2019;32(18):3109–14. doi:10.1080/14767058.2018.1455183.
236. Gürol A, Polat S. The effects of baby massage on attachment between mother and their infants. *Asian Nurs Res.* 2012;6(1):35–41. doi:10.1016/j.anr.2012.02.006.
237. Keller MK, Rehm M. A qualitative case study: fathers experiences massaging their infants. *J Behav Soc Sci.* 2015;2:29–39.
238. Improving early childhood development: WHO guideline. Geneva: World Health Organization; 2020.
239. Acceptable medical reasons for use of breast-milk substitutes. Geneva: World Health Organization, United Nations Children’s Fund; 2009.
240. Protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services: implementing the revised Baby-friendly Hospital Initiative Geneva: World Health Organization, United Nations Children’s Fund; 2018.
241. International code of marketing of breast-milk substitutes. Geneva: World Health Organization; 1981.
242. Jones E, Stewart F, Taylor B, Davis PG, Brown SJ. Early postnatal discharge from hospital for healthy mothers and term infants. *Cochrane Database Syst Rev.* 2021;(6):CD002958.
243. Yonemoto N, Nagai S, Mori R. Schedules for home visits in the early postpartum period. *Cochrane Database Syst Rev.* 2021;(7):CD009326.
244. Tiruneh GT, Shiferaw CB, Worku A. Effectiveness and cost-effectiveness of home-based postpartum care on neonatal mortality and exclusive breastfeeding practice in low-and-middle-income countries: a systematic review and meta-analysis. *BMC Pregnancy Childbirth.* 2019;19(1):507. doi:10.1186/s12884-019-2651-6.
245. Bernstein HH, Spino C, Baker A, Slora EJ, Touloukian CL, McCormick MC. Postpartum discharge: do varying perceptions of readiness impact health outcomes? *Ambul Pediatr.* 2002;2(5):388–95. doi:10.1367/1539-4409(2002)002<0388:pddvpo>2.0.co;2.
246. Dol J, Kohi T, Campbell-Yeo M, Tomblin Murphy G, Aston M, Mselle L. Exploring maternal postnatal newborn care postnatal discharge education in Dar es Salaam, Tanzania: barriers, facilitators and opportunities. *Midwifery.* 2019;77:137–43. doi:10.1016/j.midw.2019.07.009.
247. Malagon-Maldonado G, Connelly CD, Bush RA. Predictors of readiness for hospital discharge after birth: building evidence for practice. *Worldviews Evid Based Nurs.* 2017;14(2):118–27. doi:10.1111/wvn.12208.
248. Tokhi M, Comrie-Thomson L, Davis J, Portela A, Chersich M, Luchters S. Involving men to improve maternal and newborn health: a systematic review of the effectiveness of interventions. *PLoS One.* 2018;13(1):e0191620. doi:10.1371/journal.pone.0191620.
249. José Santos S. MenCare in Latin America: challenging harmful masculine norms and promoting positive changes in men’s caregiving – EMERGE Case Study 5. Brighton: Institute of Development Studies, Promundo-US, Sonke Gender Justice; 2015.
250. WHO recommendations on health promotion interventions for maternal and newborn health. Geneva: World Health Organization; 2015.
251. International Initiative for Impact Evaluation (3ie). An evidence map of social, behavioural and community engagement interventions for reproductive, maternal, newborn and child health. Geneva: World Health Organization; 2017.

252. Ruane-McAteer E, Amin A, Hanratty J, Lynn F, Corbijn van Willenswaard K, Reid E, et al. Interventions addressing men, masculinities and gender equality in sexual and reproductive health and rights: an evidence and gap map and systematic review of reviews. *BMJ Glob Health*. 2019;4(5):e001634. doi:10.1136/bmjgh-2019-001634.
253. Gunn J, Lumley J, Chondros P, Young D. Does an early postnatal check-up improve maternal health: results from a randomised trial in Australian general practice. *Br J Obstet Gynaecol*. 1998;105(9):991-7. doi:10.1111/j.1471-0528.1998.tb10263.x.
254. Timing of first postnatal contact by midwife. In: Postnatal care NICE guideline NG194. National Institute for Health and Care Excellence; 2021.
255. Dol J, Richardson B, Bonet M, Langlois EV, Parker R, Scott H, et al. Timing of maternal and neonatal mortality and morbidity in healthy women and newborns during the postnatal period: a systematic review protocol. *JBI Evid Synth*. 2021;19(3):629-43. doi:10.11124/jbies-20-00036.
256. Oza S, Cousens SN, Lawn JE. Estimation of daily risk of neonatal death, including the day of birth, in 186 countries in 2013: a vital-registration and modelling-based study. *Lancet Glob Health*. 2014;2(11):e635-44. doi:10.1016/s2214-109x(14)70309-2.
257. Oza S, Lawn JE, Hogan DR, Mathers C, Cousens SN. Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000-2013. *Bull World Health Organ*. 2015;93(1):19-28. doi:10.2471/BLT.14.139790.
258. Jones E, Taylor B, MacArthur C, Bradshaw S, Hope L, Cummins C. Early postnatal discharge for infants: a meta-analysis. *Pediatrics*. 2020;146(3):e20193365. doi:10.1542/peds.2019-3365.
259. Sainz Bueno JA, Romano MR, Teruel RG, Benjumea AG, Palacín AF, González CA, et al. Early discharge from obstetrics-pediatrics at the Hospital de Valme, with domiciliary follow-up. *Am J Obstet Gynecol*. 2005;193(3):714-26. doi:10.1016/j.ajog.2005.01.015.
260. Petrou S, Boulvain M, Simon J, Maricot P, Borst F, Perneger T, et al. Home-based care after a shortened hospital stay versus hospital-based care postpartum: an economic evaluation. *BJOG*. 2004;111(8):800-6. doi:10.1111/j.1471-0528.2004.00173.x.
261. Bowers J, Cheyne H. Reducing the length of postnatal hospital stay: implications for cost and quality of care. *BMC Health Serv Res*. 2016;16(1):16. doi:10.1186/s12913-015-1214-4.
262. Benova L, Owolabi O, Radovich E, Wong KLM, Macleod D, Langlois EV, et al. Provision of postpartum care to women giving birth in health facilities in sub-Saharan Africa: a cross-sectional study using Demographic and Health Survey data from 33 countries. *PLoS Med*. 2019;16(10):e1002943. doi:10.1371/journal.pmed.1002943.
263. Doctor HV, Radovich E, Benova L. Time trends in facility-based and private-sector childbirth care: analysis of Demographic and Health Surveys from 25 sub-Saharan African countries from 2000 to 2016. *J Glob Health*. 2019;9(2):020406. doi:10.7189/jogh.09.020406.
264. Montagu D, Sudhinaraset M, Diamond-Smith N, Campbell O, Gabrysch S, Freedman L, et al. Where women go to deliver: understanding the changing landscape of childbirth in Africa and Asia. *Health Policy Plan*. 2017;32(8):1146-52. doi:10.1093/heapol/czx060.
265. Brooten D, Roncoli M, Finkler S, Arnold L, Cohen A, Mennuti M. A randomized trial of early hospital discharge and home follow-up of women having cesarean birth. *Obstet Gynecol*. 1994;84(5):832-8.
266. Altuntuğ K, Ege E. Effects of health education on mothers' readiness for postpartum discharge from hospital, on postpartum complaints, and quality of life. *Turkish J Res Devel Nurs*. 2013;15(2):45-56.
267. Bryanton J, Beck CT, Montelpare W. Postnatal parental education for optimizing infant general health and parent-infant relationships. *Cochrane Database Syst Rev*. 2013;(11):CD004068.

268. Rasaily R, Saxena N, Pandey S, Garg BS, Swain S, Iyengar SD, et al. Effect of home-based newborn care on neonatal and infant mortality: a cluster randomised trial in India. *BMJ Glob Health*. 2020;5(9):e000680. doi:10.1136/bmjgh-2017-000680.
269. LeFevre AE, Shillcutt SD, Waters HR, Haider S, El Arifeen S, Mannan I, et al. Economic evaluation of neonatal care packages in a cluster-randomized controlled trial in Sylhet, Bangladesh. *Bull World Health Organ*. 2013;91(10):736–45. doi:10.2471/BLT.12.117127.
270. Pitt C, Tawiah T, Soremekun S, ten Asbroek AH, Manu A, Tawiah-Agyemang C, et al. Cost and cost-effectiveness of newborn home visits: findings from the Newhints cluster-randomised controlled trial in rural Ghana. *Lancet Glob Health*. 2016;4(1):e45–56. doi:10.1016/S2214-109X(15)00207-7.
271. Daviaud E, Owen H, Pitt C, Kerber K, Bianchi Jassir F, Barger D, et al. Overview, methods and results of multi-country community-based maternal and newborn care economic analysis. *Health Policy Plan*. 2017;32(suppl 1):i6–20. doi:10.1093/heapol/czx055.
272. Barger D, Owen H, Pitt C, Kerber K, Sitrin D, Mayora C, et al. Multi-country analysis of the cost of community health workers kits and commodities for community-based maternal and newborn care. *Health Policy Plan*. 2017;32(suppl 1):i84–92. doi:10.1093/heapol/czx038.
273. United Nations Children’s Fund (UNICEF), World Health Organization, World Bank Group, United Nations Inter-agency Group for Child Mortality Estimation. Levels and trends in child mortality: report 2019. New York (NY): UNICEF; 2019.
274. Chao F, You D, Pedersen J, Hug L, Alkema L. National and regional under-5 mortality rate by economic status for low-income and middle-income countries: a systematic assessment. *Lancet Glob Health*. 2018;6(5):e535–47. doi:10.1016/S2214-109X(18)30059-7.
275. McPherson R, Hodgins S. Postnatal home visitation: lessons from country programs operating at scale. *J Glob Health*. 2018;8(1):010422. doi:10.7189/jogh.08.010422.
276. Escobar GJ, Braveman PA, Ackerson L, Odouli R, Coleman-Phox K, Capra AM, et al. A randomized comparison of home visits and hospital-based group follow-up visits after early postpartum discharge. *Pediatrics*. 2001;108(3):719–27. doi:10.1542/peds.108.3.719.
277. Lieu TA, Braveman PA, Escobar GJ, Fischer AF, Jensvold NG, Capra AM. A randomized comparison of home and clinic follow-up visits after early postpartum hospital discharge. *Pediatrics*. 2000;105(5):1058–65. doi:10.1542/peds.105.5.1058.
278. WHO recommendations: optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting. Geneva: World Health Organization; 2012.
279. Guidelines for malaria vector control. Geneva: World Health Organization; 2019.
280. WHO guideline on health workforce development, attraction, recruitment and retention in rural and remote areas. Geneva: World Health Organization; 2021.
281. Baguiya A, Portela A, Moyvisan A, Gerlach N, Gopal P, Sauvé C, et al. Effectiveness of male involvement intervention on maternal and newborn health outcomes (in preparation).
282. Kunene B, Beksinska M, Zondi S, Mthembu N, Mullick S, Ottolenghi E, et al. Involving men in maternity care: South Africa. Washington (DC): Population Council; 2004.
283. WHO recommendations on home-based records for maternal, newborn and child health. Geneva: World Health Organization; 2018.
284. WHO guideline: recommendations on digital interventions for health system strengthening. Geneva: World Health Organization; 2019.