

Appendix D. Results of Statistical Models Testing for Interactions Between the Intervention and the Level of Patient-Reported Shared Decision-Making at Baseline

a. Primary Outcomes		Change in EQ5D: Quality of Life (Increase = improvement)			Change in Pain Interference (Increase=Worsening Symptoms)			Change in Sleep Disruption (range: 32-84.2; 84.2=maximum disruption)			Lower Extremity Functioning (range: 16.5-58.6; 58.6=best functioning)			Depressive Symptoms (range:38.2-81.3; 81.3=worst symptoms)		
Predictors		95%		Estimate [standard error (se)]	Confidence Interval (C.I.)	P-	Estimate (se)	P-	Estimate (se)	P-	Estimate (se)	P-	Estimate (se)	P-		
		Estimate	Value													
	Intercept	0.01 (0.01)	0.353	(2.77)	0.92	0.022	(0.53)	1.57	0.329	(2.58)	4.71	0.889	(2.57)	8.12	0.232	
	Intervention effect	0.01 (0.01)	0.380	(3.86)	4.90	0.491	(0.67)	1.52	0.751	(3.31)	6.17	0.925	(3.20)	0.96	0.097	
Shared Decision Making (SDM): Range: 0-100; higher score indicates highest level of shared decision making	SDM 1st quartile - low level (0.00 - 24.44) ^a	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
	SDM 2nd quartile (26.67 - 48.89)	0.01 (0.01)	-0.01, 0.04	0.278	0.48 (3.22)	-5.83, 6.80	0.880	(0.62)	1.12	0.883	(3.10)	4.04	0.512	(3.20)	0.75	0.084
	SDM 3rd quartile (51.11 - 73.33)	0.02 (0.01)	-0.01, 0.04	0.167	0.48 (3.09)	-7.23, 4.88	0.704	-0.3 (0.65)	1.25	0.967	(3.24)	4.06	0.479	(3.10)	0.58	0.076
	SDM 4th quartile - high level (75.56 - 100)	0.02 (0.01)	0.04	0.068	(2.99)	-7.23, 4.49	0.646	(0.59)	1.10	0.922	(2.88)	4.26	0.410	(3.02)	-5.50	-11.41,
Interaction Between Intervention and SDM	Intervention*SDM 1st quartile ^a	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
	Intervention*SDM 2nd quartile	-0.001 (0.02)	-0.04, 0.03	0.732	3.31 (4.60)	12.31	0.472	(0.86)	2.11	0.618	(4.14)	7.19	0.825	(4.11)	11.57	0.391
	Intervention*SDM 3rd quartile	-0.02 (0.02)	-0.05, 0.01	0.219	5.31 (4.37)	3.25, 13.87	0.224	(0.83)	1.41	0.803	(4.29)	9.69	0.763	(4.07)	13.57	0.169
	Intervention*SDM 4th quartile	-0.02 (0.02)	-0.05, 0.01	0.210	2.47 (4.42)	11.14	0.577	(0.78)	1.77	0.759	(3.84)	8.80	0.742	(3.80)	13.64	-0.103

a=reference group for comparison

b. Secondary Outcomes		Change in Communication (Higher Score=Better Communication)			Probability of Receiving Minimum Effective Dose 12 Month Post Treatment Start		
Predictors		95%					
		Estimate	Confidence		Estimate	P-	
		[standard error (se)]	Interval (C.I.)	P-value	(se)	95% C.I.	Value
	Intercept	0.87 (0.66)	-0.42, 2.15	0.187	0.16 (0.24)	-0.31, 0.62	0.506
	Intervention effect	-0.43 (0.97)	-2.33, 1.48	0.660	0.02 (0.34)	-0.64, 0.68	0.956
Shared Decision Making (SDM): Range: 0-100; higher score indicates highest level of shared decision making	SDM 1st quartile - low level (0.00 - 24.44) ^a	--	--	--	--	--	--
	SDM 2nd quartile (26.67 - 48.89)	-0.38 (0.77)	-1.88, 1.13	0.625	-0.15 (0.29)	-0.72, 0.43	0.619
	SDM 3rd quartile (51.11 - 73.33)	-1.27 (0.74)	-2.72, 0.19	0.089	-0.15 (0.29)	-0.71, 0.42	0.614
	SDM 4th quartile - high level (75.56 - 100)	-1.25 (0.69)	-2.61, 0.10	0.070	0.11 (0.27)	-0.63, 0.41	0.677
Interaction Between Intervention and SDM	Intervention*SDM 1st quartile ^a	--	--	--	--	--	--
	Intervention*SDM 2nd quartile	-0.35 (1.16)	-2.62, 1.93	0.766	0.16 (0.42)	-0.67, 0.99	0.701
	Intervention*SDM 3rd quartile	-0.14 (1.08)	-2.25, 1.98	0.901	0.16 (0.40)	-0.63, 0.95	0.695
	Intervention*SDM 4th quartile	0.31 (1.03)	-1.71, 2.33	0.764	0.09 (0.39)	-0.68, 0.86	0.819