

Table A2. Vulnerable Population: Descriptive Statistics of Secondary Outcomes at 30 Days and 90 Days by Study Arm

	N	Vulnerable Population		
	N	Usual Care N=185	GUIDED-HF N=183	P-value
KCCQ (baseline)	360	46.61 (32.29, 68.49)	50.78 (37.63, 65.62)	0.408
KCCQ (30 days)	268	54.43 (35.42, 74.61)	66.15 (46.88, 80.73)	0.006
KCCQ (changes in 30 days)	263	5.469 (-7.422, 17.06)	10.76 (-1.562, 23.18)	0.067
KCCQ (90 days)	266	61.46 (39.58, 80.21)	63.54 (44.92, 83.33)	0.421
KCCQ (changes in 90 days)	260	9.375 (-2.604, 24.74)	11.98 (-4.167, 28.12)	0.706
Dutch (baseline)	368	10 (8, 11)	11 (9, 12)	0.061
Dutch (30 days)	368	9 (0, 11)	10 (0, 12)	0.002
Dutch (changes in 30 days)	368	0 (-7, 1)	0 (-3, 1)	0.12
Dutch (90 days)	366	8 (0, 11)	11 (0, 12)	<0.001
Dutch (changes in 90 days)	366	-1 (-8, 1)	0 (-2, 2)	0.009
PROMIS Anxiety (30 days)	265	49.4 (37.1, 57.4)	49.4 (37.1, 58.4)	0.528
PROMIS Anxiety (90 days)	256	45.9 (37.1, 56.4)	45.9 (37.1, 57.4)	0.488
PROMIS Depression (30 days)	265	47.5 (38.2, 56.57)	47.5 (38.2, 55.1)	0.749
PROMIS Depression (90 days)	258	47.5 (38.2, 55.9)	44.7 (38.2, 55.1)	0.688
ED Discharge Satisfaction	312			0.507
Strongly Disagree	312	1% (2)	0% (0)	
Disagree	312	1% (1)	1% (1)	
Neither Agree nor Disagree	312	4% (6)	3% (5)	
Agree	312	46% (72)	41% (64)	
Strongly Agree	312	48% (75)	55% (86)	

Note: Values are proportions (frequencies) for categorical variables, median (IQR: lower quartile, upper quartile) for continuous variables.