

Table 1c.ii - Comparisons of subject characteristics in imputed data – MOST

| Variable | TKR (n=154) | Non-TKR (n=154) | TKR minus non-TKR <i>Delta (Δ)</i> and [95% CI] | Effect Size |
|---|----------------|--------------------|--|-------------|
| <i>mean +/- standard deviation (SD)</i> | | | | (Δ/SD) |
| Age | 65.49 ± 7.18 | 65.21 ± 7.18 | 0.28 [-1.32 , 1.88] | 0.02 |
| Male, N(%) | 0.31 ± 0.46 | 0.31 ± 0.46 | 0.01 [-0.10 , 0.11] | 0.01 |
| Baseline BMI | 32.33 ± 9.48 | 31.44 ± 7.18 | 0.89 [-0.98 , 2.77] | 0.05 |
| Baseline SF-12 Physical | 36.80 ± 7.83 | 38.62 ± 10.43 | -1.82 [-3.87 , 0.24] | -0.1 |
| Baseline SF-12 Mental | 55.99 ± 11.00 | 54.23 ± 11.80 | 1.76 [-0.79 , 4.31] | 0.08 |
| Baseline WOMAC Knee pain (0-100) | 40.23 ± 24.82 | 35.33 ± 22.24 | 4.90 [-0.36 , 10.17] | 0.1 |
| Baseline Knee Pain in Contralateral (0-100) | 21.46 ± 20.06 | 19.38 ± 24.10 | 2.08 [-2.87 , 7.03] | 0.05 |
| Baseline Hip Pain or Pain/Ache/Stiff | 0.46 ± 0.50 | 0.61 ± 0.50 | -0.15 [-0.26 , -0.04] | -0.15 |
| At least one comorbidity, N (%) | 0.38 ± 0.49 | 0.39 ± 0.49 | -0.01 [-0.12 , 0.10] | -0.01 |
| Narcotics, N (%) | 0.20 ± 0.44 | 0.13 ± 0.39 | 0.07 [-0.03 , 0.16] | 0.08 |
| Follow-up SF-12 Physical | 40.01 ± 12.11 | 40.10 ± 11.99 | -0.09 [-2.78 , 2.60] | 0 |
| Follow-up WOMAC Knee Pain (0-100) | 14.01 ± 17.26 | 23.93 ± 17.26 | -9.92 [-13.77 , -6.06] | -0.29 |

*Shaded rows indicate variables where definitions varied between databases so that these variables ultimately were excluded as candidates in the building of final models.