

Table S12. Treatment preferences among men by whether or not a respondent had experienced discrimination by a medical provider

utility	Odds Ratio	Standard Error	z	p	95% CI	
Always reliable	1.889	0.115	10.420	<0.001	1.676	2.128
Always understandable language	1.681	0.122	7.180	<0.001	1.459	1.937
Talk therapy only	0.694	0.093	-2.740	0.006	0.535	0.902
Both medication and talk therapy	1.671	0.127	6.780	<0.001	1.441	1.939
30 minute commute	0.860	0.058	-2.230	0.026	0.754	0.982
60 minute commute	0.364	0.035	-10.560	<0.001	0.301	0.439
\$20 cost	0.725	0.053	-4.400	<0.001	0.628	0.837
\$75 cost	0.208	0.022	-14.720	<0.001	0.169	0.256
Always reliable*Discrimination	1.129	0.222	0.620	0.536	0.768	1.660
Always understandable language*Discrimination	0.956	0.154	-0.280	0.779	0.697	1.310
Talk therapy only*Discrimination	1.620	0.487	1.600	0.109	0.899	2.922
Both medication and talk therapy*Discrimination	1.242	0.244	1.100	0.270	0.845	1.824
30 minute commute*Discrimination	1.321	0.215	1.710	0.088	0.959	1.818
60 minute commute*Discrimination	1.608	0.378	2.020	0.044	1.014	2.550
\$20 cost*Discrimination	0.837	0.127	-1.170	0.240	0.621	1.127
\$75 cost*Discrimination	0.803	0.160	-1.100	0.269	0.544	1.186