

Table S13. Treatment preferences among women by whether or not a respondent had experienced discrimination by a medical provider

utility	Odds Ratio	Standard Error	z	p	95% CI	
Always reliable	1.971	0.126	10.650	<0.001	1.740	2.233
Always understandable language	1.892	0.121	10.010	<0.001	1.670	2.144
Talk therapy only	1.055	0.124	0.450	0.651	0.837	1.329
Both medication and talk therapy	2.180	0.175	9.720	<0.001	1.863	2.551
30 minute commute	0.910	0.053	-1.610	0.107	0.812	1.021
60 minute commute	0.400	0.028	-12.930	<0.001	0.348	0.460
\$20 cost	0.603	0.039	-7.760	<0.001	0.531	0.685
\$75 cost	0.172	0.015	-11.400	<0.001	0.145	0.205
Always reliable*Discrimination	0.786	0.104	-1.820	0.069	0.606	1.019
Always understandable language*Discrimination	0.970	0.145	-0.200	0.840	0.723	1.301
Talk therapy only*Discrimination	0.692	0.169	-1.510	0.131	0.429	1.116
Both medication and talk therapy*Discrimination	1.002	0.164	0.010	0.992	0.727	1.380
30 minute commute*Discrimination	1.001	0.118	0.010	0.994	0.794	1.262
60 minute commute*Discrimination	0.893	0.152	-0.660	0.506	0.640	1.246
\$20 cost*Discrimination	1.061	0.128	0.490	0.624	0.837	1.344
\$75 cost*Discrimination	1.279	0.268	1.170	0.241	0.848	1.929