

Table S16. Treatment preferences among men by whether or not a respondent had experienced discrimination by both a medical provider and front desk staff

utility	Odds Ratio	<i>Standard Error</i>	z	p	95% CI	
Always reliable	1.923	0.116	10.800	<0.001	1.708	2.165
Always understandable language	1.652	0.115	7.240	<0.001	1.442	1.893
Talk therapy only	0.720	0.092	-2.560	0.010	0.560	0.926
Both medication and talk therapy	1.700	0.124	7.290	<0.001	1.474	1.960
30 minute commute	0.869	0.057	-2.140	0.032	0.764	0.988
60 minute commute	0.360	0.034	-10.850	<0.001	0.299	0.433
\$20 cost	0.715	0.050	-4.820	<0.001	0.624	0.820
\$75 cost	0.208	0.021	-15.570	<0.001	0.171	0.253
Always reliable*Discrimination	1.022	0.255	0.090	0.930	0.627	1.666
Always understandable language*Discrimination	1.112	0.214	0.550	0.581	0.762	1.623
Talk therapy only*Discrimination	1.617	0.594	1.310	0.191	0.787	3.323
Both medication and talk therapy*Discrimination	1.259	0.319	0.910	0.365	0.765	2.069
30 minute commute*Discrimination	1.426	0.259	1.950	0.051	0.998	2.036
60 minute commute*Discrimination	2.303	0.447	4.300	<0.001	1.574	3.369
\$20 cost*Discrimination	0.828	0.164	-0.950	0.343	0.561	1.222
\$75 cost*Discrimination	0.683	0.162	-1.610	0.107	0.430	1.086