

Table S3. Treatment preferences among non-Hispanic white respondents by whether or not a respondent had experienced discrimination by a medical provider

utility	Odds Ratio	Standard error	z	p	95% CI	
Always reliable	2.056	0.118	12.530	<0.001	1.837	2.302
Always understandable language	1.918	0.120	10.370	<0.001	1.696	2.169
Talk therapy only	0.798	0.093	-1.930	0.053	0.635	1.003
Both medication and talk therapy	1.876	0.136	8.660	<0.001	1.627	2.163
30 minute commute	0.869	0.049	-2.480	0.013	0.778	0.971
60 minute commute	0.353	0.026	-13.960	<0.001	0.305	0.409
\$20 cost	0.617	0.040	-7.490	<0.001	0.544	0.700
\$75 cost	0.168	0.014	-21.620	<0.001	0.143	0.198
Always reliable*Discrimination	0.832	0.154	-1.000	0.320	0.579	1.195
Always understandable language*Discrimination	0.845	0.145	-0.990	0.324	0.604	1.181
Talk therapy only*Discrimination	0.936	0.317	-0.200	0.844	0.481	1.819
Both medication and talk therapy*Discrimination	1.655	0.304	2.740	0.006	1.155	2.372
30 minute commute*Discrimination	1.063	0.197	0.330	0.743	0.739	1.527
60 minute commute*Discrimination	0.794	0.194	-0.940	0.345	0.491	1.283
\$20 cost*Discrimination	0.909	0.125	-0.690	0.488	0.694	1.190
\$75 cost*Discrimination	0.952	0.200	-0.240	0.813	0.631	1.436