

Table A2: Comparative treatment effects. Effect sizes reflect between-group pairwise comparisons adjusted for covariates in the final mixed effects general linear model (95% confidence interval). Rows labeled “treatment” and “6-month” show the post-treatment and follow-up scores relative to pre-treatment scores, respectively. A positive between-group difference for the MAL means that the group listed first in the comparison showed greater gains in arm use. A negative between-group difference for the WMFT means that the group listed first in the comparison showed greater gains. Statistically significant contrasts are indicated with an *. Clinically meaningful differences between groups are italicized.

	Gaming+ vs CIMT	Gaming vs CIMT	Gaming+ vs Standard Care	Gaming vs Standard Care	Gaming+ vs Gaming	CIMT vs Standard Care
MAL treatment	-0.2 (-0.4, 0.1)	-0.5 (-0.8, -0.3)*	0.9 (0.6, 1.2)	0.5 (0.2, 0.8)*	0.4 (0.1, 0.7)*	1.2 (0.8, 1.5)*
MAL 6-month	-0.3 (-0.6, 0.0)	-0.6 (-0.8, -0.3)*	0.4 (0.1, 0.8)*	0.1 (-0.3, 0.4)	0.3 (0.0, 0.7)	0.7 (0.4, 1.1)*
WMFT treatment	0.11 (-0.03, 0.24)	<i>0.20</i> <i>(0.06, 0.33)*</i>	-0.10 (-0.24, 0.04)	-0.01 (-0.15, 0.13)	-0.09 (-0.22, 0.04)	<i>-0.21 (-0.35, -0.07)*</i>
WMFT 6-month	0.10 (-0.10, 0.30)	0.16 (-0.04, 0.36)	0.01 (-0.19, 0.21)	0.07 (-0.13, 0.27)	-0.06 (-0.25, 0.14)	-0.09 (-0.30, 0.12)