







**Figure 52: Sensitivity and specificity of diarrhoea for diagnosing adrenal insufficiency (cortisol peak response < 11 µg after low dose 10 µg short corticotropin stimulation test) in people with HIV-infection**

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Casanova-Cardiel 2003	8	13	39	46	0.17 [0.08, 0.31]	0.78 [0.65, 0.88]		



**Figure 53: Sensitivity and specificity of diarrhoea for diagnosing adrenal insufficiency (twofold value of basal cortisol after low dose 10 µg short corticotropin stimulation test) in people with HIV-infection**

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Casanova-Cardiel 2003	17	4	61	24	0.22 [0.13, 0.33]	0.86 [0.67, 0.96]		

**Figure 54: Sensitivity and specificity of diarrhoea for diagnosing adrenal insufficiency (any cortisol value > 18 µg/dL after low dose 10 µg short corticotropin stimulation test) in people with HIV-infection**

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Casanova-Cardiel 2003	0	21	5	80	0.00 [0.00, 0.52]	0.79 [0.70, 0.87]		

**Figure 55: Sensitivity and specificity of diarrhoea for diagnosing adrenal insufficiency (any cortisol value > 20 µg/dL after low dose 10 µg short corticotropin stimulation test) in people with HIV-infection**

Study	TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)
Casanova-Cardiel 2003	0	21	6	79	0.00 [0.00, 0.46]	0.79 [0.70, 0.87]		

**Figure 56: Sensitivity and specificity of diarrhoea for diagnosing adrenal hypofunction (all cortisol measurements <18 µg/dL including 1 µg/mL ACTH test)**

