Figure 12: Sensitivity and specificity of mucoses hyperpigmentation for diagnosing adrenal insufficiency (cortisol peak response < 11 μg after low dose 10 μg short corticotropin stimulation test) in people with HIV-infection

Study	ТΡ	FP	FΝ	ΤN	Sensitivity (95% CI)	Specificity (95% CI) Sensitivity (95% C	(I) Specificity (95% CI)
Casanova-Cardiel 2003	3	4	44	55	0.06 [0.01, 0.18]	0.93 (0.84,0.98)	

Figure 13: Sensitivity and specificity of mucoses hyperpigmentation for diagnosing adrenal insufficiency (twofold value of basal cortisol after low dose 10 µg short corticotropin stimulation test) in people with HIV-infection

Study	ΤP	FP	FN	ΤN	Sensitivity (95% CI)	Specificity (95% CI) Sensitivity (95% CI)	Specificity (95% CI)
Casanova-Cardiel 2003	3	4	75	24	0.04 [0.01, 0.11]	0.86 [0.67,0.96] <mark>- </mark>	

Figure 14: Sensitivity and specificity of mucoses hyperpigmentation for diagnosing adrenal insufficiency (any cortisol value > 18 μg/dL after low dose 10 μg short corticotropin stimulation test) in people with HIV-infection

Study	ТΡ	FP	FN	ΤN	Sensitivity (95% CI)	Specificity (95% CI) Sensitivity (95% CI) Specificity (95% CI)
Casanova-Cardiel 2003	1	6	4	95	0.20 [0.01, 0.72]	

Figure 15: Sensitivity and specificity of mucoses hyperpigmentation for diagnosing adrenal insufficiency (any cortisol value > 20 μg/dL after low dose 10 μg short corticotropin stimulation test) in people with HIV-infection

Study
TP
FP
FN
TN
Sensitivity (95% CI)
Specificity (95% CI)