

NLM Citation: Qian H. GABA_C Receptors in the Vertebrate Retina. 2005 May 1 [Updated 2007 Jun 1]. In: Kolb H, Fernandez E, Jones B, et al., editors. Webvision: The Organization of the Retina and Visual System [Internet]. Salt Lake City (UT): University of Utah Health Sciences Center; 1995-. **Bookshelf URL:** https://www.ncbi.nlm.nih.gov/books/

GABA_C Receptors in the Vertebrate Retina

WEBVISION The Organization of the Rectina and Visual System Map the Astronometer of Paper New

Haohua Qian

Created: May 1, 2005; Updated: June 1, 2007.

Properties of GABA Receptors

 γ -Aminobutyric acid (GABA) is the main inhibitory neurotransmitter in the central nervous system. The inhibitory action of GABA is mediated by the receptors present on the cell membrane and results in a reduction of neuronal excitability. At least three types of GABA receptors have been characterized. Table 1 summarizes some of the general properties of these three types of GABA receptors. GABAA receptors are ligand-gated chloride channels. They mediate fast inhibition and have a wide distribution throughout the central nervous system. GABA_A receptors have a diverse molecular composition. At least 16 subunits in six groups have been identified. Pharmacologically, these receptors are antagonized by bicuculline. GABAA receptors are also the targets of many therapeutic compounds (such as general anaesthetics, sedative drugs, and alcohols). These compounds allosterically modulate GABAA receptor channel activities. GABAB receptors belong to the Gprotein-coupled receptor superfamily. The inhibition of GABAB receptors is mediated by indirect gating of either potassium or calcium channels. GABAB receptors are activated by baclofen and antagonized by phaclofen and saclofen. The subunits of GABAB receptors have recently been cloned. GABAC receptors are the newly identified member of the GABA receptor family. They are also linked to chloride channels, with distinct physiological and pharmacological properties. In contrast to the fast and transient responses elicited from GABA_A receptors, GABA_C receptors mediate slow and sustained responses. Pharmacologically, GABA_C receptors are bicuculline- and baclofen-insensitive and are not modulated by many GABAA receptor modulators (such as benzodiazepines and barbiturates). GABA ρ subunits are thought to participate in forming GABA_C receptors on the neuronal membrane, but the exact molecular composition of these receptors is yet to be determined. GABA_C receptors are expressed in many brain regions, with prominent distributions on retinal neurons, suggesting that these receptors play important roles in retinal signal processing.

	GABA _A receptor	GABA _B receptor	$GABA_C$ receptor
Category	Ligand-gated channel	G-protein-coupled receptor	Ligand-gated channel
Subunits	$\alpha,\beta,\gamma,\delta,\epsilon,\pi$	GBR1, GBR2	ρ
Agonists	Muscimol, THIP	Baclofen	
Antagonists	Bicuculline, picrotoxin	Phaclofen	TPMPA, picrotoxin
Desensitization	Yes	No	No

Table 1. Characteristics of GABA receptors.

Table 1 continued from previous page.

	GABA _A receptor	GABA _B receptor	GABA _C receptor
Modulator	Benzodiazepine barbiturates		Zinc

GABA_C Responses on Retinal Neurons

The term "GABA_C receptor" was first used by Johnston to describe a novel GABA binding site on neuronal membranes (1, 2). Although recent studies indicate a wide distribution of GABA_C receptors in many parts of the central nervous system (3-7), these receptors are most prominently expressed in the vertebrate retina. In the fish white perch retina, rod-driven (H4) horizontal cells were the first retinal neurons where GABA_C receptors were characterized (8). Subsquently, GABA_C receptor-mediated responses have been detected in many types of retinal neurons, including bipolar cells (9-15), cone-driven horizontal cells in catfish (16, 17), cone photoreceptors (18), and ganglion cells (19). Among all these retinal neurons, the rod-driven horizontal cells of white perch are the only cells where GABA responses are mediated solely by GABA_C receptors. The GABA responses elicited from other cells are usually a mixture of GABA receptors and/or GABA transporters. The unique properties of the white perch rod-driven horizontal cell provided an excellent model to characterize the physiological and pharmacological properties of GABA_C receptors on retinal neurons (8, 20).

An example of a solitary rod-driven horizontal cell isolated from white perch retina is shown in Fig. 1, together with a typical GABA-elicited response from such a cell. These horizontal cells receive input from rod photoreceptors in the retina; and when isolated, they keep their typical morphology in culture. Rod-driven horizontal cells have a flat cell body with diameter of 50-100 µm. There are several thick primary dendrites from which many fine processes extend. As shown in Fig. 1, GABA elicits a slow and sustained response from these cells. The GABA-induced membrane currents are mediated by chloride ions and, therefore, exhibit inhibitory actions on these neurons. The responses showed no sign of desensitization, i.e., the responses are maintained at a steady level as long as GABA is present. Furthermore, the GABA responses elicited from these horizontal cells exhibit slow kinetics, which could best be observed in the offset response. After the termination of GABA application, the membrane current returns to the baseline very slowly, with a time constant of ~15 seconds. Such slow and sustained response properties are typical of GABA_C receptors.

It is interesting to note that the neurons in distal retina (i.e., photoreceptors, horizontal cells, and bipolar cells) do not produce action potentials at all. They only generate slow graded responses to light stimuli. It has always been thought that retinal neurons must have special ways to process and analyze such slow signals compared with fast transient neurons of the brain. The kinetics of GABA_C receptor-mediated responses are thus particularly suited for the generatation of signals in distal retinal neurons.

Pharmacology of GABA_C Receptors

GABA_C receptors exhibit a distinct pharmacology that differs from classical GABA_A or GABA_B receptors. GABA_C receptors were first described by Johnston for bicuculline- and baclofen-insensitive GABA binding sites on neuronal membranes (1, 2). More detailed studies indicated that GABA_C receptors on retinal neurons are not sensitive to the competitive antagonists of either GABA_A receptors (such as SR95531 and hydrastine) or GABA_B receptors (such as phaclofen and saclofen). Because the competitive antagonists are thought to interact with the GABA binding sites on the receptors, these results indicate that a different conformation of GABA molecule is preferred for binding to the GABA_C receptor. In agreement with such a notion, the specific agonists of GABA_A and GABA_B receptors exhibit quite different activity on GABA_C receptors. They either have no effect (isonipecotic acid, baclofen), act as partial agonists (isoguvacine, muscimol), or as antagonists (THIP, P4S, 3-APA, and 3-APMPA) (20, 21). I4AA, a partial agonist of GABA_A receptors, behaves as a potent antagonist on



Figure 1. An example of a rod-driven (H4) horizontal cell (left) isolated from white perch retina. GABA elicits a sustained and slow response (right) from these cells mediated by GABA_C receptors. Thus far, this is the only preparation in which the GABA response is mediated solely by GABA_C receptors.

GABA_C responses of retinal neurons and as a partial agonist on expressed GABA_C receptors in *Xenopus* oocytes (20, 22).

GABA_C receptors also differ from classical GABA_A receptors in terms of their responses to various modulators. Two groups of compounds, benzodiazepine and barbiturates, are well-known to modulate GABA_A activity. On the other hand, none of these compounds have any significant influence on responses mediated by GABA_C receptors (23). For example, the GABA-elicited response on white perch rod-driven horizontal cells are virtually identical in the presence or absence of either diazepam or pentobarbital (8). Another class of GABA_A receptor modulators, known as neuroactive steroids, has different effects on the expressed GABA_C receptor in *Xenopus* oocytes. Whereas some of them modulate GABA responses on these receptors, others do not (24, 25). The effect of these neuroactive steroids on GABA_C receptors on neurons has yet to be determined.

Although both GABA_A and GABA_C receptors are linked to chloride channels, the channel properties of these two receptors are quite different. Unlike GABA_A receptors, picrotoxin inhibition on GABA_C receptors in white perch horizontal cells exhibits both competitive and non-competitive mechanisms (20). In mammalian retina (rat), on the other hand, GABA_C receptors are insensitive to picrotoxin blockage (9, 13). The unusual features of GABA_C receptors in rat retina are attributed to a single amino acid substitution in the receptor subunit (26). Furthermore, TBPS (like picrotoxin, another chloride channel blocker on GABA_A receptors) does not block the responses mediated by GABA_C receptors (20, 27). In addition, GABA_C receptor-gated chloride channels exhibit a very small single channel conductance (14, 28).

GABA_C receptor activities are modulated by divalent cations (17, 29-31). In particular, GABA_C receptormediated responses are inhibited by low concentrations of zinc ions. The high sensitivity of GABA_C receptors to zinc inhibition is attributed to a histidine residue on the extracellular domain of the subunits (32). Recently, a new GABA_C receptor antagonist, TPMPA, has become available commercially. This compound is thought to be a specific inhibitor of the GABA_C receptor (33). The availability of such a drug will greatly facilitate further studies of this receptor.

Molecular Biology of GABAC Receptors

The GABA_C receptor is a member of the ligand-gated channel superfamily. By analogy to the well-studied nicotinic acytocholine receptors, GABA_C receptors are thought to exhibit the structure shown schematically in Fig. 2. These receptors are pentamers, i.e., five subunits constitute the functional channel (34). The receptors have a long extracellular domain containing ligand binding sites and several modulatory sites. In the middle of the receptor, GABA gates an ionic channel. Binding of GABA to the receptor induces a conformational change in receptor structure, which leads to the opening of the channel.

For each subunit forming the receptor, the structure is thought to be that shown in Fig. 3. The subunit contains a long extracellular N-terminal domain that has ligand binding sites, four transmembrane domains, and a large intracellular loop that connects the third and the fourth transmembrane domains. The ionic channel is formed by the second transmembrane domain of each subunit. For the large intracellular loop, there are several putative phosphorylation sites, indicated as colored circles with letter "P" in the figure. Phosphorylation of these residues is implicated for modulation of receptor activities. For example, it has been reported that dopamine modulates GABA_C receptor activity in both catfish cone-driven horizontal cells and tiger salamander bipolar cell terminals (35, 36). On rat bipolar cells, the GABA_C receptor activities are modulated by protein kinase C (37). The modulation of receptor activities by intracellular second messenger systems is also observed at GABA_C receptors expressed on *Xenopus* oocytes (38). However, the mechanisms for modulation by intracellular second messager systems are yet to be determined. Recently, Filippova and coauthors (39) provided evidence for GABA_C receptor internalization upon phosphorylation of the subunits. In addition, there is evidence that the large intracellular loop of GABA_C receptor subunits is involved in the interaction of receptor protein with other intracellular proteins, which may play an important role in the clustering of the receptors on neuronal membranes (40).

There is much evidence to indicate that GABA_C receptors are composed of GABA ρ subunits. These were first cloned from a human retinal cDNA library (41, 42). When expressed in *Xenopus* oocytes, GABA ρ subunits formed functional homo-oligomeric receptors with properties similar to those of GABA_C receptors in retinal neurons (43). Furthermore, the expression of GABA ρ subunits has been detected on retinal neurons where GABA_C receptor-mediated responses have been recorded (44-46). In white perch retina, we have now cloned five forms of GABA ρ subunits (22, 46). Fig. 4 shows a comparison of white perch GABA ρ subunits and those cloned from mammalian retinas. The distances between the various connecting elements represent the degree of divergence among subunits. Unlike the mammalian retina, where only one form of $\rho 1$ and $\rho 2$ subunits has been identified, in white perch there are two forms of the subunit for each p1 and p2 family. In accordance with their deduced amino acid sequences and the properties of the receptors they formed on *Xenopus* oocytes, each $\rho 1$ and ρ^2 family was subdivided into A and B forms. All white perch GABA ρ^1 and ρ^2 subunits are able to form functional homo-oligometric receptors when expressed in *Xenopus* oocytes. GABA-elicited responses in these expressed receptors are sustained, bicuculline insensitive, and are not modulated by either benzodiazepines or barbiturates, features typical of GABAC receptors. Similar to GABAC receptors on retinal neurons, GABA p receptors also gate chloride channels. However, the receptors expressed by each of the GABA ρ subunits display unique response properties that distinguish one from the other. For example, the sensitivity of GABA activation and picrotoxin inhibition varies among subunits. In addition, I4AA acts as an antagonist on A-type ρ receptors, whereas it is a partial agonist on B-type ρ receptors (22).

The waveform of receptor-mediated responses plays an important role in shaping the neuronal signal. Interestingly, the kinetics of the GABA response are also different for the receptors formed by each individual subunits. Current-trace responses to application of 10 μ M GABA are illustrated in Fig. 5. These recordings

indicate that there are significant differences in the kinetics of the GABA responses from *Xenopus* oocytes, depending on which white perch GABA ρ subunitis are expressed. To quantitate the kinetics of the GABA response, the offset GABA responses (current traces after GABA application is terminated) were replotted on a semi-logarithmic scale, with the amplitudes normalized to their initial values (Fig. 6). In each case, the data were fit by a straight line, indicating that offset responses can be described by a single exponential function. The slope of the line represents the time constant of the decay and shows that the receptors formed by the various p subunits exhibit significant differences in their response kinetics. The average time constants of offset responses elicited from receptors formed by various white perch GABA ρ subunits are shown in the bar graph in Fig. 6C. There are consistent differences between the response kinetics of the two receptor families and between their subgroups. For example, the responses from $\rho 1$ receptors were significantly slower than those of $\rho 2$ receptors. Such difference in the response kinetics among p1 and p2 receptors are determined, in large part, by a single residue at the second transmembrane domain of the subunits (47). This dichotomy among $\rho 1$ and $\rho 2$ subunits is well conserved in all species where GABA ρ subunits have been cloned. $\rho 1$ subunits, which have a proline at the residue site, combine to make a receptor with slower kinetics, whereas ρ^2 subunits, which contain a serine at the residue site, form receptors with faster response rate. Thus, receptors made of human ρ 1 subunits exhibit slower response kinetics than receptors made of $\rho 2$ subunits (5). The kinetic differences among the receptors formed by various GABA ρ subunits could provide building blocks for the nevous system to construct different types of signal filters and different types of neuronal signaling in the nervous system.

Function of GABA_C Receptors in the Retina

Although rod-driven horizontal cells provide an excellent model in which to characterize GABA_C receptors in the retina, recent studies indicate that GABA_C receptors are present on various other types of retinal neurons. GABA_C receptor-mediated responses have been recorded from cone-driven horizontal cells in catfish (16, 17), cone photoreceptors (18), and some types of ganglion cells (19). GABA_C responses are particularly prominent in bipolar cells of every species examined thus far (9-12, 14, 15), and both immunocytochemistry and *in situ* hybridization studies indicate that GABA_C receptors are present on bipolar cells (15, 44, 45, 48). It appears that these receptors play an important role in shaping signal transmission from bipolar cells to third-order neurons in the retina.

Fig. 7 illustrates some examples of bipolar cells isolated from white perch retina. These bipolar cells keep their morphology when isolated in culture. They usually have a pear-shaped cell body from which several dendrites and one axon extend. The GABA responses of bipolar cells in white perch retina have both transient and sustained components, indicating both GABA_A and GABA_C receptors are present, as shown in Fig. 8. The transient component can be selectively blocked by the co-application of bicuculline, leaving a more sustained response. Thus, the electrophysiological and pharmacological properties of GABA_C receptors on bipolar cells are very similar to those of GABA_C receptors on rod-driven horizontal cells (11, 14, 37, 49).

Different kinetic properties of GABA_A and GABA_C receptors suggest that they play different roles in mediating inhibition on bipolar cell terminals (15, 46, 50). Furthermore, various subtypes of bipolar cells exhibit different proportions of GABA_A and GABA_C receptors. For example, in the rat retina, there is a clear difference in the contribution of GABA_A and GABA_C receptors to rod and cone bipolar cells (51). In white perch, too, different morphological types of bipolar cells exhibit different proportions of GABA_C receptor-mediated components (14). These results strongly suggest that different subtypes of bipolar cell use various mixtures of GABA_A and GABA_C receptors to perform different activities and help create the variety of functional pathways through the retina.

Because of the presence of multiple GABA receptors on retinal neurons, it is sometimes difficult to isolate the contributions of each receptor. Recent studies on ganglion cell responses reveal some interesting features of GABA_C receptors in retinal information processing. For example, activation of GABA_C receptors leads to more



Figure 2. Schematic diagram of GABA_C receptors. These receptors are formed by five subunits with an ionic channel in the middle of the receptor. On the extracellular side, the receptor contains the binding site for GABA and several modulatory sites.

transient light responses in ganglion cells (52) and the delayed inhibition mediated by GABA_C receptors is thought to play a major role in shaping edge-enhancement of ganglion cell receptive fields (53). The bipolar cell to ganglion cell synapse is probably heavily influenced by inhibitory amacrine feed-forward or feedback synapses, and these appear to be via primarily GABA_C receptors.



Figure 3. Schematic digram of the GABA_C receptor subunits. Each subunit contains a long extracellular N-terminal region that contains two cystine residues thought to form disulfide bonds. The extracellular domain forms the GABA binding sites and other modulatory sites. The subunit crosses the cell membrane four times with a short C-terminal region on the outside of the cell. The second transmembrane domain forms the ionic channel of the receptor. There are several putative phosphorylation sites on the large intracellular loop between the third and the fourth transmembrane domains (colored circles with letter "P").



Figure 4. Dendrogram showing the relation of the GABA ρ subunits cloned from white perch retina to those cloned from mammalian retinas. Analysis was performed on the deduced amino acid sequence of each subunit.



Figure 5. GABA (10 μ M) induced responses from white perch GABA ρ subunits expressed in *Xenopus* oocytes. The duration of GABA application is shown by the bar above each trace. Variations in response amplitude often result from the different level of receptor expression in individual oocytes; response kinetics are a relatively stable feature, independent of amplitude.



Figure 6. Kinetics of the GABA responses obtained from the white perch GABA ρ receptors expressed in *Xenopus* oocytes. A. Semilogarithmic plot of GABA responses measured at various times following the offset of drug application (10 μ M GABA); the responses were normalized to their initial amplitude at the time of offset. The responses are well fit by straight lines, indicating that the relationship can be described by a single exponential function. B. A similar analysis applied to the offset GABA responses obtained from perch ρ 2 receptors. Note different time scales in A and B. C. Comparison of the time constants calculated from GABA offset responses elicited from various GABA ρ receptors.



Figure 7. Examples of solitary white perch bipolar cells in culture.



Figure 8. GABA-elicited responses from isolated white perch bipolar cells. The membrane currents elicited by GABA application (left) contain both transient and sustained components, indicating both GABA_A and GABA_C receptors are present. In the pesence of bicuculline, GABA_A receptor activity is inhibited, revealing sustained GABA responses mediated by GABA_C receptors (right).

About the Author



Dr. Haohua Qian was born in Jiangsu, China. He received his B.A. in Biology from Nanjing University (1982), M.S. in Neurobiology from Shanghai Institute of Physiology (1985), and Ph.D. in Anatomy and Cell Biology from University of Illinois at Chicago (1991). He is currently an Assistant Professor of Neurosciences in the Department of Ophthalmology and Visual Sciences at the University of Illinois at Chicago. During his postdoctoral studies with Dr. John E. Dowling at Harvard University, he characterized a new type of GABA receptor, the GABA_C ρ receptor, on retinal neurons. He is currently continuing these studies on the molecular structure and physiological functions of GABA_C receptors in the vertebrate retina.

References

- 1. Johnston GAR. Multiplicity of GABA receptors. In: Olsen RW, Venter JC, editors. Receptor biochemistry and methodology. Vol. 5. Alan R. Liss, Inc.; 1986. p. 57-71.
- 2. Johnston GAR. GABAc receptors: relatively simple transmitter-gated ion channels? Trends Pharmacol Sci. 1996;17:319–323. PubMed PMID: 8885697.
- 3. Albrecht BE, Breitenbach U, Stuhmer T, Harvey RJ, Darlison MG. In situ hybridization and reverse transcription-polymerase chain reaction studies on the expression of the GABA(C) receptor rho1- and rho2-subunit genes in avian and rat brain. Eur J Neurosci. 1997;9:2414–2422. PubMed PMID: 9464935.
- 4. Boue-Grabot E, Roudbaraki M, Bascles L, Tramu G, Bloch B, Garret M. Expression of GABA receptor rho subunits in rat brain. J Neurochem. 1998;70:899–907. PubMed PMID: 9489708.
- 5. Enz R, Cutting GR. GABAC receptor rho subunits are heterogeneously expressed in the human CNS and form homo- and heterooligomers with distinct physical properties. Eur J Neurosci. 1999;11:41–50. PubMed PMID: 9987010.
- 6. Sivilotti L, Nistri A. GABA receptor mechanisms in the central nervous system. Prog Neurobiol. 1991;36:35–92. PubMed PMID: 1847747.
- 7. Wegelius K, Pasternack M, Hiltunen JO, Rivera C, Kaila K, Saarma M, Reeben M. Distribution of GABA receptor rho subunit transcripts in the rat brain. Eur J Neurosci. 1998;10:350–357. PubMed PMID: 9753143.
- 8. Qian H, Dowling JE. Novel GABA responses from rod-driven retinal horizontal cells. Nature. 1993;361:162–164. PubMed PMID: 8421521.
- 9. Feigenspan A, Wässle H, Bormann J. Pharmacology of GABA receptor Cl⁻ channels in rat retinal bipolar cells. Nature. 1993;361:159–163. PubMed PMID: 7678450.
- 10. Lukasiewicz PD, Wong ROL. GABAC receptors on ferret retinal bipolar cells: a diversity of subtypes in mammals? Vis Neurosci. 1997;14:989–994. PubMed PMID: 9364734.
- 11. Lukasiewicz PD, Maple BR, Werblin FS. A novel GABA receptor on bipolar cell terminals in the tiger salamander retina. J Neurosci. 1994;14:1202–1212. PubMed PMID: 8120620.
- 12. Nelson R, Schaffner AE, Li Y-X, Walton MC. Distribution of GABA(C)-like responses among acutely dissociated rat retinal neurons. Vis Neurosci. 1999;16:179–190. PubMed PMID: 10022489.
- 13. Pan Z-H, Lipton SA. Multiple GABA receptor subtypes mediate inhibition of calcium influx at rat retinal bipolar cell terminals. J Neurosci. 1995;15:2668–2679. PubMed PMID: 7722621.
- 14. Qian H, Dowling JE. GABAA and GABAC receptors on hybrid bass retinal bipolar cells. J Neurophysiol. 1995;74:1920–1928. PubMed PMID: 8592185.
- 15. Qian H, Li L, Chappell RL, Ripps H. GABA receptors of bipolar cells from the skate retina: actions of zinc on GABA-mediated membrane currents. J Neurophysiol. 1997b;78:2402–2412. PubMed PMID: 9356392.
- 16. Dong CJ, Picaud SA, Werblin FS. GABA transporters and GABAC-like receptors on catfish cone- but not rod-driven horizontal cells. J Neurosci. 1994;14:2648–2658. PubMed PMID: 8182433.
- 17. Kaneda M, Mochizuki M, Kaneko A. Modulation of GABAC response by Ca2+ and other divalent cations in horizontal cells of the catfish retina. J Gen Physiol. 1997;110:741–747. PubMed PMID: 9382900.
- Picaud S, Pattnaik B, Hicks D, Forster V, Fontaine V, Sahel J, Dreyfus H. GABAA and GABAC receptors in adult porcine cones: evidence from a photoreceptor-glia co-culture model. J Physiol. 1998;513:33–42. PubMed PMID: 9782157.
- 19. Zhang J, Slaughter MM. Perferential suppression of the ON pathway by GABAC receptors in the amphibian retina. J Neurophysiol. 1995;74:1583–1592. PubMed PMID: 8989395.
- 20. Qian H, Dowling JE. Pharmacology of novel GABA receptors found on rod horizontal cells of the white perch retina. J Neurosci. 1994;14:4299–4307. PubMed PMID: 8027780.
- 21. Woodward RM, Polenzani L, Miledi R. Characterization of bicuculline/baclofen-insensitive (r-like) gaminobutyric acid receptors expressed in Xenopus oocytes. II. pharmacology of g-aminobutyric acid A and g-aminobutyric acid B receptor agonists and antagonists. Mol Pharmacol. 1993;43:609–625. PubMed PMID: 8386310.

- 22. Qian H, Dowling JE, Ripps H. Molecular and pharmacological properties of GABA-rho subunits from white perch retina. J Neurobiol. 1998;37:305–320. PubMed PMID: 9805275.
- 23. Polenzani L, Woodward RM, Miledi R. Expression of mammalian g-aminobutyric acid receptors with distinct pharmacology in Xenopus oocytes. Proc Natl Acad Sci U S A. 1991;88:4318–4322. PubMed PMID: 1709741.
- 24. Morris KD, Moorefield CN, Amin J. Differential modulation of the gamma-aminobutyric acid type C receptor by neuro active steroids. Mol Pharmacol. 1999;56:752–759. PubMed PMID: 10496958.
- 25. Woodward RM, Polenzani L, Miledi R. Effects of steroids on g-aminobutyric acid receptors expressed in Xenopusoocytes by poly(A)+ RNA from mammalian brain and retina. Mol Pharmacol. 1992a;41:89–103. PubMed PMID: 1370710.
- 26. Zhang D, Pan ZH, Zhang X, Brideau AD, Lipton SA. Cloning of a gamma-aminobutyric acid type C receptor subunit in rat retina with a methionine residue critical for picrotoxin in channel block. Proc Natl Acad Sci U S A. 1995;92:11756–11760. PubMed PMID: 8524843.
- 27. Woodward RM, Polenzani L, Miledi R. Characterization of bicuculline/baclofen-insensitive g-aminobutyric acid receptors expressed in Xenopus oocytes I. effects of Cl- channel inhibitors. Mol Pharmacol. 1992b;42:165–173. PubMed PMID: 1378924.
- 28. Chang Y, Weiss DS. Channel opening locks agonist on to the GABAC receptor. Nat Neurosci. 1999;2:219–225. PubMed PMID: 10195213.
- 29. Calvo DJ, Vazquez AE, Miledi R. Cationic modulation of r1-type g-aminobutyrate receptors expressed in Xenopus oocytes. Proc Natl Acad Sci U S A. 1994;91:12725–12729. PubMed PMID: 7809110.
- 30. Chang Y, Amin J, Weiss DS. Zinc is a mixed antagonist of homomeric r1 g-aminobutyric acid-activated channels. Mol Pharmacol. 1995;47:595–602. PubMed PMID: 7535381.
- 31. Dong CJ, Werblin FS. Zinc down modulates the GABAc receptor current in cone horizontal cells acutely isolated from the catfish retina. J Neurophysiol. 1995;73:916–919. PubMed PMID: 7760150.
- 32. Wang TL, Hackam A, Guggino WB, Cutting GR. A single histidine residue is essential for zinc inhibition of GABA rho 1 receptors. J Neurosci. 1995;15:7684–7691. PubMed PMID: 7472519.
- 33. Ragozzino D, Woodward RM, Murata Y, Eusebi F, Overman LE, Miledi R. Design and in vitro pharmacology of a selective gamma-aminobutyric acid C receptor antagonist. Mol Pharmacol. 1996;50:1024–1030. PubMed PMID: 8863850.
- 34. Amin J, Weiss DS. Insights into the activation mechanism of rho1 GABA receptors obtained by coexpression of wild type and activation-impaired subunits. Proc Biol Sci. 1996;263:273–282. PubMed PMID: 8920250.
- 35. Dong CJ, Werblin FS. Dopamine modulation of GABAC receptor function in an isolated retinal neuron. J Neurophysiol. 1994;71:1258–1260. PubMed PMID: 8201417.
- Wellis DP, Werblin FS. Dopamine modulates GABAc receptors mediating inhibition of calcium entry into and transmitter release from bipolar cell terminals in tiger salamander retina. J Neurosci. 1995;15:4748– 4761. PubMed PMID: 7623108.
- 37. Feigenspan A, Bormann J. Modulation of GABAC receptors in rat retinal bipolar cells by protein kinase C. J Physiol. 1994a;481:325–330. PubMed PMID: 7738828.
- 38. Kusama T, Sakurai M, Kizawa Y, Uhl GR, Murakami H. GABA rho1 receptor: inhibition by protein kinase C activators. Eur J Pharmacol. 1995;291:431–434. PubMed PMID: 8719430.
- 39. Filippova N, Dudley R, Weiss DS. Evidence for phosphorylation-dependent internalization of recombinant human r1 GABAC receptors. J Physiol. 1999;518:385–399. PubMed PMID: 10381587.
- 40. Hanley JG, Koulen P, Bedford F, Gordon-Weeks PR, Moss SJ. The protein MAP-1B links GABA(C) receptors to the cytoskeleton at retinal synapses. Nature. 1999;397:66–69. PubMed PMID: 9892354.
- 41. Cutting GR, Lu L, Zoghbi H, O'Hara BF, Kasch LM, Montrose-Rafizader C, Donovan DM, Shimada S, Antonarakis SE, Guggino W, Uhl GR, Kazazian HH. Cloning of the g-aminobutyric acid (GABA) rho1 cDNA: a GABA receptor subunit highly expressed in the retina. Proc Natl Acad Sci U S A. 1991;88:2673–2677. PubMed PMID: 1849271.

- 42. Cutting GR, Curristin S, Zoghbi H, O'Hara B, Selden MF, Uhl GR. Identification of a putative gammaaminobutyric acid (GABA) receptor subunit rho2 cDNA and colocalization of the genes encoding rho2 (GABRR2) and rho1 (GABRR1) to human chromosome 6q14-q21 and mouse chromosome 4. Genomics. 1992;12:801–806. PubMed PMID: 1315307.
- 43. Shimada S, Cutting G, Uhl GR. g-Aminobutyric acid A or C receptor? g-Aminobutyric acid r1 receptor RNA induces bicuculline-, barbiturate-, and benzodiazepine-insensitive g-aminobutyric acid responses in Xenopus oocytes. Mol Pharmacol. 1992;41:683–687. PubMed PMID: 1314944.
- 44. Enz R, Brandstätter JH, Hartveit E, Wässle H, Bormann J. Expression of GABA receptor rho 1 and rho2 subunits in the retina and brain of the rat. Eur J Neurosci. 1995;7:1495–1501. PubMed PMID: 7551175.
- 45. Enz R, Brandstätter JH, Wässle H, Bormann J. Immunocytochemical localization of the GABAC receptor rho subunits in the mammalian retina. J Neurosci. 1996;16:4479–4490. PubMed PMID: 8699258.
- 46. Qian H, Hyatt G, Schanzer A, Hazra R, Hackam A, Cutting GR, Dowling JE. A comparison of GABAC and rho subunit receptors from the white perch retina. Vis Neurosci. 1997a;14:843–851. PubMed PMID: 9364723.
- 47. Qian H, Dowling JE, Ripps H. A single amino acid in the second transmembrane domain of GABA r subunits is a determinant of the response kinetics of GABAC receptors. J Neurobiol. 1999;40:67–76. PubMed PMID: 10398072.
- 48. Koulen P, Brandstätter JH, Kröger S, Enz R, Bormann J, Wässle H. Immunocytochemical localization of the GABA(C) receptor rho subunits in the cat, goldfish, and chicken retina. J Comp Neurol. 1997;380:520–532. PubMed PMID: 9087530.
- 49. Feigenspan A, Bormann J. Differential pharmacology of GABAA and GABAC receptors on rat retinal bipolar cells. Eur J Pharmacol. 1994b;288:97–104. PubMed PMID: 7535710.
- 50. Lukasiewicz PD, Shields CR. Different combinations of GABAA and GABAC receptors confer distinct temporal properties to retinal synaptic responses. J Neurophysiol. 1998;79:3157–3167. PubMed PMID: 9636116.
- 51. Euler T, Wassle H. Different contributions of GABAA and GABAC receptors to rod and cone bipolar cells in a rat retinal slice preparation. J Neurophysiol. 1998;79:1384–1395. PubMed PMID: 9497419.
- 52. Dong CJ, Werblin FS. Temporal contrast enhancement via GABAC feedback at bipolar terminals in the tiger salamander retina. J Neurophysiol. 1998;79:2171–2180. PubMed PMID: 9535976.
- 53. Jacobs AL, Werblin FS. Spatio temporal pattern sat the retinal output. J Neurophysiol. 1998;80:447–451. PubMed PMID: 9658062.

License

All copyright for chapters belongs to the individual authors who created them. However, for non-commercial, academic purposes, images and content from the chapters portion of Webvision may be used with a non-exclusive rights under a Attribution, Noncommercial 4.0 International (CC BY-NC) Creative Commons license. Cite Webvision, http://webvision.med.utah.edu/ as the source. Commercial applications need to obtain license permission from the administrator of Webvision and are generally declined unless the copyright owner can/wants to donate or license material. Use online should be accompanied by a link back to the original source of the material. All imagery or content associated with blog posts belong to the authors of said posts, except where otherwise noted.