




## Luliconazole

Revised: June 21, 2021.

CASRN: 187164-19-8



## Drug Levels and Effects

### Summary of Use during Lactation

Topical luliconazole has not been studied during breastfeeding. Because it is poorly absorbed after topical use and is highly plasma protein bound, it is a low risk to the nursing infant. Avoid application to the nipple area and ensure that the infant's skin does not come into direct contact with the areas of skin that have been treated. Only water-miscible cream or gel products should be applied to the breast because ointments may expose the infant to high levels of mineral paraffins via licking.[1]

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

*Infant Levels.* Relevant published information was not found as of the revision date.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## References

1. Noti A, Grob K, Biedermann M, et al. Exposure of babies to C(15)-C(45) mineral paraffins from human milk and breast salves. *Regul Toxicol Pharmacol.* 2003;38:317–25. PubMed PMID: 14623482.

## Substance Identification

### Substance Name

Luliconazole

### CAS Registry Number

187164-19-8

### Drug Class

Breast Feeding

Lactation

Antifungal Agents

Anti-Infective Agents