




Sodium Ferric Gluconate Complex

Revised: September 15, 2023.

CASRN: 34089-81-1



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of sodium ferric gluconate complex during breastfeeding. Ferrlecit contains 9 mg/mL of benzyl alcohol, which may enter breastmilk and be absorbed by the infant. An alternate intravenous drug with more published data available may be preferred. For additional information on iron use during breastfeeding, see the monograph on [Iron Salts](#).

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Ferric Carboxymaltose, Ferric Derisomaltose, Iron Sucrose

Substance Identification

Substance Name

Sodium Ferric Gluconate Complex

CAS Registry Number

34089-81-1

Drug Class

Breast Feeding

Lactation

Milk, Human

Ferric Compounds

Hematinics

Iron Compounds