




## Edrophonium

Revised: December 3, 2018.

CASRN: 312-48-1



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of edrophonium during breastfeeding. However, because of its short half-life and quaternary ammonium structure, it is unlikely to be excreted into breastmilk or orally absorbed by the infant. Administering the drug just after breastfeeding and waiting 2 to 3 hours before breastfeeding again should avoid any adverse reactions in the infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Edrophonium

### CAS Registry Number

312-48-1

### Drug Class

Breast Feeding

Lactation

Cholinesterase Inhibitors

Diagnostic Agents

Parasympathomimetics