




## Peramivir

Revised: February 7, 2019.

CASRN: 330600-85-6



## Drug Levels and Effects

### Summary of Use during Lactation

Because peramivir is poorly absorbed orally, it is not likely to reach the bloodstream of the infant in clinically important amounts. However, because no information is available on the use of peramivir breastfeeding, an alternate drug may be preferred, especially while nursing a newborn or preterm infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Oseltamivir, Zanamivir

## Substance Identification

### Substance Name

Peramivir

### CAS Registry Number

330600-85-6

### Drug Class

Breast Feeding

Lactation

Antiviral Agents

Neuraminidase Inhibitors