




Ixazomib

Revised: December 3, 2018.

CASRN: 1072833-77-2



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of ixazomib during breastfeeding. Because its half-life is about 9.5 days, it is likely to accumulate in the infant. It is also given in combination with leflunomide and dexamethasone, which may increase the risk to the infant. The manufacturer recommends that breastfeeding be discontinued during ixazomib therapy and for 90 days after the last dose.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Ixazomib

CAS Registry Number

1072833-77-2

Drug Class

Breast Feeding

Lactation

Antineoplastic Agents

Enzyme Inhibitors

Protein Kinase Inhibitors

Signal Transduction Inhibitors