




Simethicone

Revised: March 17, 2021.

CASRN: 8050-81-5



Drug Levels and Effects

Summary of Use during Lactation

Although no data are available on the use of simethicone during breastfeeding, it is known that simethicone is not absorbed orally. Therefore, it cannot be transferred to breastmilk. It is also used safely in breastfed infants. [1,2] No special precautions are required.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

References

1. Hagemann TM. Gastrointestinal medications and breastfeeding. *J Hum Lact.* 1998;14:259–62. PubMed PMID: 10205441.
2. Savino F, Pelle E, Palumeri E, et al. *Lactobacillus reuteri* (American Type Culture Collection Strain 55730) versus simethicone in the treatment of infantile colic: A prospective randomized study. *Pediatrics.* 2007;119:e124–30. PubMed PMID: 17200238.

Substance Identification

Substance Name

Simethicone

CAS Registry Number

8050-81-5

Drug Class

Breast Feeding

Lactation

Antifoaming Agents