


U.S. National Library of Medicine National Center for Biotechnology Information **NLM Citation:** Drugs and Lactation Database (LactMed®) [Internet]. Bethesda (MD): National Institute of Child Health and Human Development; 2006-. Frovatriptan. [Updated 2024 Apr 15]. **Bookshelf URL:** https://www.ncbi.nlm.nih.gov/books/



# Frovatriptan

Revised: April 15, 2024.

CASRN: 158747-02-5



# **Drug Levels and Effects**

## Summary of Use during Lactation

There is no published experience with frovatriptan during breastfeeding. If frovatriptan is required by the mother of an older infant, it is not a reason to discontinue breastfeeding, but until more data become available, an alternate drug may be preferred, especially while nursing a newborn or preterm infant.[1] Painful, burning nipples and breast pain have been reported after doses of sumatriptan and other triptans. This has occasionally been accompanied by a decrease in milk production.

### **Drug Levels**

Maternal Levels. Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Infant Levels. Relevant published information was not found as of the revision date.

### **Effects in Breastfed Infants**

Relevant published information was not found as of the revision date.

#### **Effects on Lactation and Breastmilk**

A review of four European adverse reaction databases found 26 reported cases of, painful, burning nipples, painful breasts, breast engorgement and/or painful milk ejection in women who took a triptan while nursing. Pain was sometimes intense and occasionally led to decreased milk production. Pain generally subsided with time as the drug was eliminated. The authors proposed that triptans may cause vasoconstriction of the arteries in the breast, nipples, and the arteries surrounding the alveoli and milk ducts, causing a painful sensation and a painful milk ejection reflex.[2]

#### **Alternate Drugs to Consider**

Eletriptan, Rizatriptan, Sumatriptan, Zolmitriptan

#### References

- 1. Amundsen S, Nordeng H, Nezvalova-Henriksen K, et al. Pharmacological treatment of migraine during pregnancy and breastfeeding. Nat Rev Neurol 2015;11:209-19. PubMed PMID: 25776823.
- 2. Conijn M, Maas V, van Tuyl M, et al. Breastfeeding-related adverse drug reactions of triptans: A descriptive analysis using four pharmacovigilance databases. Breastfeed Med 2024. PubMed PMID: 38563407.

## **Substance Identification**

#### **Substance Name**

Frovatriptan

### **CAS Registry Number**

158747-02-5

#### **Drug Class**

Breast Feeding Lactation Milk, Human Serotonin Receptor Agonists Serotonin 5-HT1 Receptor Agonists Triptans Vasoconstrictor Agents