




Telmisartan

Revised: January 16, 2023.

CASRN: 144701-48-4



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of telmisartan during breastfeeding and the manufacturer advises avoidance of breastfeeding during telmisartan therapy. An alternate drug may be preferred, especially while nursing a newborn or preterm infant.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Candesartan, Captopril, Benazepril, Enalapril, Quinapril

Substance Identification

Substance Name

Telmisartan

CAS Registry Number

144701-48-4

Drug Class

Breast Feeding

Lactation

Milk, Human

Antihypertensive Agents

Angiotensin II Type 1 Receptor Blockers

Angiotensin Receptor Blockers

ARBs