




## Articaïne

Revised: December 3, 2018.

CASRN: 23964-58-1



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of articaïne during breastfeeding. Based on the low excretion of other local anesthetics into breastmilk, a single dose of articaïne injected during breastfeeding is unlikely to adversely affect the breastfed infant. However, an alternate drug may be preferred, especially while nursing a newborn or preterm infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Articaine

### CAS Registry Number

23964-58-1

### Drug Class

Breast Feeding

Lactation

Anesthetics, Local