




Estazolam

Revised: February 15, 2023.

CASRN: 29975-16-4



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of estazolam during breastfeeding. Because of the long duration of action of estazolam, an alternate hypnotic is preferred, especially while nursing a newborn or preterm infant. If estazolam is used, monitor the infant for sedation, poor feeding and poor weight gain.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Zaleplon, Zolpidem

Substance Identification

Substance Name

Estazolam

CAS Registry Number

29975-16-4

Drug Class

Breast Feeding

Lactation

Milk, Human

Benzodiazepines

Hypnotics and Sedatives