




## Castor

Revised: June 21, 2021.

CASRN: 8001-79-4



## Drug Levels and Effects

### Summary of Use during Lactation

Castor (*Ricinus communis*) beans contain triglycerides, mostly consisting of ricinoleic acid esters, and small amounts of the toxic ricin and ricine. Pressing of the beans produces castor oil and purification of the oil eliminates the ricin and ricine. Castor oil is a strong stimulant laxative. Castor beans as well as a homeopathic preparation of castor purportedly reduce milk flow,[1] but it is also reportedly used as a galactagogue.[2-4] A poultice of castor leaves is a purported galactagogue.[5,6] In some parts of India, castor oil is also reportedly applied to the breasts to stimulate lactation.[6] No scientifically valid clinical trials support any of these uses and some preparations may be toxic to the infant. Galactagogues should never replace evaluation and counseling on modifiable factors that affect milk production.[7,8] No data exist on the excretion of any components of the castor plant or castor oil into breastmilk or on their safety and efficacy in nursing mothers or infants. However,

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

little of the active ricinoleic acid is thought to be absorbed from the intestine. Because of a lack of information, other cathartics may be preferred in nursing mothers.

In traditional Indian culture, castor oil has been administered to newborn infants during the first 2 to 3 days of life, often resulting in adverse effects.[9,10] Administration of castor oil to newborns is dangerous and should be avoided.

Dietary supplements do not require extensive pre-marketing approval from the U.S. Food and Drug Administration. Manufacturers are responsible to ensure the safety, but do not need to *prove* the safety and effectiveness of dietary supplements before they are marketed. Dietary supplements may contain multiple ingredients, and differences are often found between labeled and actual ingredients or their amounts. A manufacturer may contract with an independent organization to verify the quality of a product or its ingredients, but that does *not* certify the safety or effectiveness of a product. Because of the above issues, clinical testing results on one product may not be applicable to other products. More detailed information [about dietary supplements](#) is available elsewhere on the LactMed Web site.

## Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

## Effects in Breastfed Infants

In rural India, castor oil has been traditionally given to infants during the first 2 to 3 days of life to clear the intestine of meconium. This practice can result in paralytic ileus and aspiration pneumonia.[9] Severe hypoalbuminemia was also reported in a 1.5-month-old infant whose grandmother gave him castor oil daily from the fifth day of life, resulting in diarrhea and malnutrition.[10]

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

(Cathartic) [Bisacodyl](#), [Magnesium Hydroxide](#), [Senna](#)

## References

1. Hardy ML. Women's health series: Herbs of special interest to women. J Am Pharm Assoc (Wash). 2000;40:234–42. PubMed PMID: 10730024.
2. Eglash A. Treatment of maternal hypergalactia. Breastfeed Med. 2014;9:423–5. PubMed PMID: 25361472.
3. Winterfeld U, Meyer Y, Panchaud A, et al. Management of deficient lactation in Switzerland and Canada: A survey of midwives' current practices. Breastfeed Med. 2012;7:317–8. PubMed PMID: 22224508.
4. Dandotiya H, Singh G, Kashaw SK. The galactagogues use by Indian tribal communities to over come poor lactation. Int J Biotechnol Bioeng Res. 2013;4:243–8.
5. Yarnell E. Botanical medicine in pregnancy and lactation. Altern Complement Ther. 1997;3(April):93–100.
6. Rasiya Beegam A, Nayar TS. Plants used for natal healthcare in folk medicine of Kerala, India. Indian J Tradit Knowl. 2011;10:523–7.
7. Brodribb W. ABM Clinical Protocol #9. Use of galactagogues in initiating or augmenting maternal milk production, second revision 2018. Breastfeed Med. 2018;13:307–14. PubMed PMID: 29902083.
8. Breastfeeding challenges: ACOG Committee Opinion, Number 820. Obstet Gynecol. 2021;137:e42–e53. PubMed PMID: 33481531.

9. Benakappa DG, Raju M. Breast-feeding practices in rural Karnataka (India) with special reference to lactation failure. *Acta Paediatr Jpn.* 1989;31:391–8. Shivananda, et al. PubMed PMID: 2514560.
10. Jayaprakash DG, Raghu Raman TS, Singh D, et al. Laxative induced hypoalbuminemia. *Indian Pediatr.* 1995;32:1037–8. PubMed PMID: 8935275.

## Substance Identification

### Substance Name

Castor

### Scientific Name

*Ricinus communis*

### CAS Registry Number

8001-79-4

### Drug Class

Breast Feeding

Lactation

Cathartics

Complementary Therapies

Oils

Phytotherapy

Plants, Medicinal