



## Factor Xa (recombinant), Inactivated

Revised: April 20, 2020.

CASRN: 1262449-58-0

Structure  
not available

## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the clinical use of coagulation factor Xa during breastfeeding. Because it is a large protein molecule with a molecular weight of about 40,000, the amount in milk is likely to be very low and absorption is unlikely because it is probably destroyed in the infant's gastrointestinal tract. Until more data become available, coagulation factor X should be used with caution during breastfeeding, especially while nursing a newborn or preterm infant.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

---

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Factor Xa (recombinant), Inactivated

### CAS Registry Number

1262449-58-0

### Drug Class

Breast Feeding

Lactation

Blood Coagulation Factors

Coagulants