| Table 9: CI | inical evidence | profile: Com | parison 4.2. | Telemedicine | versus usual care |
|-------------|-----------------|--------------|--------------|--------------|-------------------|
|-------------|-----------------|--------------|--------------|--------------|-------------------|

| Quality assessment | | | | | | No of patients | | Effect | | | | |
|----------------------|---|--|---|-------------------|-----------------|-------------------------------------|------------------|---------------|-----------------------------|--------------|---------------|------------|
| No of studi es | Design | Risk of bias | Inconsistenc y | Indirectnes s | Imprecisio n | Other consideration s | Telemedicin e | Usual care | Relati ve (95% CI) | Absolu te | Qual ity | Importance |
| Chang | Change in quality of life- CFQOL body (Follow-up: 6 months; range of scores: 0-100; Better indicated by lower values) | | | | | | | | | | | |
| 1 (Wilki | al studies | very no serious seriou inconsistenc s ¹ y | no serious Not indirectnes calculable s | Not calculable | none ulable | 4 | 3 | 3 - | Not estima ble | VER Y | IMPORTAN T | |
| nson 2008) | | | | | | Significant improvemen t at 6 | | | | LOW | | |

 $\ensuremath{\textcircled{\sc online \sc on$

| Quality assessment | | | | | | No of patients | | Effect | | | | |
|----------------------|--------|-----------------|-------------------|------------------|-----------------|-----------------------------|-------------------|---------------|-----------------------------|--------------|-------------|------------|
| No of studi es | Design | Risk of bias | Inconsistenc y | Indirectnes s | Imprecisio n | Other consideration s | Telemedicin e | Usual care | Relati ve (95% CI) | Absolu te | Qual ity | Importance |
| | | | | | | | months, p=0.02 | | | | | |

Abbreviations: CI: confidence interval; CFQOL: cystic fibrosis quality of life questionnaire 1 The quality of the evidence was downgraded by 2 because of incomplete reporting and high-loss to follow-up