




Pasireotide

Revised: November 30, 2022.

CASRN: 396091-73-9



Drug Levels and Effects

Summary of Use during Lactation

The excretion of pasireotide into breastmilk has not been studied. However, because it has a high molecular weight of 1047 daltons it is likely to be poorly excreted into breastmilk and it is a peptide that is likely digested in the infant's gastrointestinal tract. It is unlikely to reach the clinically important levels in infant serum. However, the manufacturer states that nursing mothers should not use pasireotide. An alternate drug is preferred.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Octreotide

Substance Identification

Substance Name

Pasireotide

CAS Registry Number

396091-73-9

Drug Class

Breastfeeding

Lactation

Milk, Human

Hormones

Somatostatin Analogs and Derivatives