41 Acronyms and abbreviations

ACSAcute coronary syndromesAESAnti-embolism stockingsBNFBritish National FormularyCEACost-effectiveness analysisCIConfidence intervalCPMContinuous passive motionCRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntersvice Care UnitIPOLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRANetwork meta-analysisNMANetwork meta-analysisNMANetwork meta-analysisNMANetwork meta-analysisNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Suppl AgencyPEPulmonary embolis pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTaua Embolic Scoring System <th>Acronym or abbreviation</th> <th>Description</th>	Acronym or abbreviation	Description
BNFBritish National FormularyCEACost-effectiveness analysisCIConfidence intervalCPMContinuous passive motionCRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHIDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboenolic pulmonary hypertensionPTSPast-thrombotic syndromeSCSubutaneousUKOSSUnited Kingdom Obstetric SyreemUKOSSUnited Kingdom Obstetric SyreemUKOSUnited Kingdom Obstetric Syreem	ACS	Acute coronary syndromes
BNFBritish National FormularyCEACost-effectiveness analysisCIConfidence intervalCPMContinuous passive motionCRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHIDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboenolic pulmonary hypertensionPTSPast-thrombotic syndromeSCSubutaneousUKOSSUnited Kingdom Obstetric SyreemUKOSSUnited Kingdom Obstetric SyreemUKOSUnited Kingdom Obstetric Syreem	AES	Anti-embolism stockings
CEACost-effectiveness analysisCIConfidence intervalCPMContinuous passive motionCRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGraduated compression stockingsGRADEGraduated compression stockingsHDHigh doseHDHigh doseHESHospital acquired thrombosis or hospital associated thrombosisHITHospital acquired thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntermittent pneumatic compression devicesISSInjury severity scoreIVLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombolic Soring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUKRSSUnited Kingdom Obstetric Surveillance System	BNF	
CIConfidence intervalCPMContinuous passive motionCRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMajor bleedingMHRANetwork meta-analysisNMANetwork meta-analysisNMESNumber needed to traatOACOral anticoagulantsPFEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombote syndremeSCSubcutaneousUKOSSUnited Kingdom Obstetric SystemUFHUnfractionated heparin	CEA	
CRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIncremental cost-effectiveness ratioICUInternittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPTSPost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	CI	
CRTCatheter related thrombosisCUACost-utility analysisCVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIncremental cost-effectiveness ratioICUInternittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPTSPost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	СРМ	Continuous passive motion
CVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUInternsittent pneumatic compression devicesISSInjury severity scoreIVInternsittent pneumatic compression devicesLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thromboembolic pulmonary hypertensionPTSPost-thromboembolic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	CRT	
CVCCentral venous cathetersDOACDirect oral anticoagulantDVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUInternsittent pneumatic compression devicesISSInjury severity scoreIVInternsittent pneumatic compression devicesLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thromboembolic pulmonary hypertensionPTSPost-thromboembolic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	CUA	Cost-utility analysis
DVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRANedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNumber needed to treatOACOral anticoagulantsPASAPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombolic scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	CVC	
DVTDeep vein thrombosisFIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICCRIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRANedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNumber needed to treatOACOral anticoagulantsPASAPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombolic scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	DOAC	Direct oral anticoagulant
FIDFoot impulse devicesGCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASAMSP verbasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	DVT	
GCSGraduated compression stockingsGRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUInternitent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASAMS Purchasing and Supply AgencyPEPulmonary embolics nydromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	FID	
GRADEGrading of Recommendations Assessment, Development and EvaluationHATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASAMHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	GCS	
HATHospital acquired thrombosis or hospital associated thrombosisHDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASAMHS Purchasing and Supply AgencyPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	GRADE	
HDHigh doseHESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPFPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	НАТ	
HESHospital Episode StatisticsHITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPFPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	HD	
HITHeparin-induced thrombocytopaeniaICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	HES	-
ICERIncremental cost-effectiveness ratioICUIntensive Care UnitIPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASAMHS Purchasing and Supply AgencyPEPulmonary embolismPTSPost-thrombotic syndromeSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	НІТ	
IPCDIntermittent pneumatic compression devicesISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	ICER	
ISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	ICU	Intensive Care Unit
ISSInjury severity scoreIVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	IPCD	Intermittent pneumatic compression devices
IVIntravenousLDLow doseLMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	ISS	
LMWHLow molecular weight heparinLOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	IV	
LOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	LD	Low dose
LOSLength of stayMBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	LMWH	Low molecular weight heparin
MBMajor bleedingMHRAMedicines and Healthcare Products Regulatory AgencyNMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSSubcutaneousSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	LOS	
NMANetwork meta-analysisNMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	MB	Major bleeding
NMESNeuromuscular electrical stimulationNNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	MHRA	Medicines and Healthcare Products Regulatory Agency
NNTNumber needed to treatOACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	NMA	Network meta-analysis
OACOral anticoagulantsPASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	NMES	Neuromuscular electrical stimulation
PASANHS Purchasing and Supply AgencyPEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	NNT	Number needed to treat
PEPulmonary embolismPHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	OAC	Oral anticoagulants
PHTChronic thromboembolic pulmonary hypertensionPTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	PASA	NHS Purchasing and Supply Agency
PTSPost-thrombotic syndromeSCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	PE	Pulmonary embolism
SCSubcutaneousTESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	PHT	Chronic thromboembolic pulmonary hypertension
TESSTrauma Embolic Scoring SystemUKOSSUnited Kingdom Obstetric Surveillance SystemUFHUnfractionated heparin	PTS	Post-thrombotic syndrome
UKOSS United Kingdom Obstetric Surveillance System UFH Unfractionated heparin	SC	Subcutaneous
UFH Unfractionated heparin	TESS	Trauma Embolic Scoring System
	UKOSS	United Kingdom Obstetric Surveillance System
VKA Vitamin K antagonist	UFH	Unfractionated heparin
	VKA	Vitamin K antagonist

© NICE 2018. All rights reserved. Subject to Notice of rights.

Acronym or abbreviation	Description
VTE	Venous thromboembolism