

C-reactive protein (<10 mg/l) at time of blood culture

Sensitivity and specificity



C-reactive protein (<10 mg/l)

C-reactive protein (<10 mg/l)



Neonatal infection: antibiotics for prevention and treatment evidence reviews for investigations before starting treatment for late-onset neonatal infection FINAL (April 2021)


Likelihood ratios

H-Blommendahl 2002 3.72 [2.13, 6.48] Boo 2008 5.16 [2.35, 11.34] Him H Boonkasidecha 2013 52.89 [3.39, 826.06] Joji 2018 1.48 [1.24, 1.76] H. Khair 2012 H#H 2.77 [1.71, 4.47] Kumar 2010 6.49 [2.77, 15.23] Mkony 2014 нфн, 2.96 [2.27, 3.85] Philip 1980 ⊢⊨– 3.21 [2.04, 5.06] Sakha 2008 2.49 [1.65, 3.75] ⊢∎⊣ Sharma 1993 1.92 [1.17, 3.14] Sucilathangam 2012 1.61 [0.80, 3.23] West 2012 2.84 [2.25, 3.57] ι÷Η Anwar ul Hap 2019 H∎-3.51 [2.34, 5.25] Balasubramanin 2018 H∎Hİ 1.70 [1.32, 2.19] Summary estimate 2.77 [2.33, 3.29] 4 0.1 0.5 4 16 64 256

C-reactive protein (<10 mg/l)

Positive likelihood ratio

C-reactive protein (<10 mg/l)



Neonatal infection: antibiotics for prevention and treatment evidence reviews for investigations before starting treatment for late-onset neonatal infection FINAL (April 2021)