

Intrapartum care

Quality standard

Published: 10 December 2015

www.nice.org.uk/guidance/qs105

Contents

Quality statements	4
Quality statement 1: Choosing birth setting.....	5
Quality statement.....	5
Rationale	5
Quality measures	5
What the quality statement means for different audiences.....	6
Source guidance.....	7
Definitions of terms used in this quality statement	7
Quality statement 2: One-to-one care	9
Quality statement.....	9
Rationale	9
Quality measures	9
What the quality statement means for different audiences.....	10
Source guidance.....	11
Definitions of terms used in this quality statement	11
Quality statement 3: Cardiotocography and initial assessment of a woman in labour	12
Quality statement 4: Stopping cardiotocography	13
Quality statement.....	13
Rationale	13
Quality measures	13
What the quality statement means for different audiences.....	14
Source guidance.....	14
Definitions of terms used in this quality statement	15
Quality statement 5: Interventions during labour	16
Quality statement.....	16
Rationale	16
Quality measures	16

What the quality statement means for different audiences.....	17
Source guidance.....	17
Definitions of terms used in this quality statement	18
Quality statement 6: Delayed cord clamping	19
Quality statement.....	19
Rationale	19
Quality measures	19
What the quality statement means for different audiences.....	20
Source guidance.....	21
Definitions of terms used in this quality statement	21
Quality statement 7: Skin-to-skin contact	22
Quality statement.....	22
Rationale	22
Quality measures	22
What the quality statement means for different audiences.....	23
Source guidance.....	23
Update information.....	24
About this quality standard.....	25
Resource impact.....	25
Diversity, equality and language	26

This standard is based on CG190, NG4 and NG201.

This standard should be read in conjunction with QS75, QS46, QS35, QS22, QS15, QS60, QS32, QS109, QS115, QS135 and QS192.

Quality statements

Statement 1 Women at low risk of complications during labour are given the choice of all 4 birth settings and information about local birth outcomes.

Statement 2 Women in established labour have one-to-one care and support from an assigned midwife.

Statement 3 This statement has been removed. For more details see [update information](#).

Statement 4 Women at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Statement 5 Women at low risk of complications are not offered amniotomy or oxytocin if labour is progressing normally.

Statement 6 Women do not have the cord clamped earlier than 1 minute after the birth unless there is concern about cord integrity or the baby's heartbeat.

Statement 7 Women have skin-to-skin contact with their babies after the birth.

Quality statement 1: Choosing birth setting

Quality statement

Women at low risk of complications during labour are given the choice of all 4 birth settings and information about local birth outcomes.

Rationale

Women at low risk of complications during labour and birth need information that is specific to their local or neighbouring area about safety and outcomes for women and babies in the different birth settings. This information will help women to make informed choices about where to have their baby.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

a) Evidence of local arrangements to provide women at low risk of complications with a choice of all 4 birth settings.

Data source: Local data collection.

b) Evidence of local arrangements to provide women at low risk of complications with local information about birth outcomes.

Data source: Local data collection.

Process

a) Proportion of women at low risk of complications with a recorded discussion at their 16-week antenatal appointment of their preferred choice of birth setting.

Numerator – The number in the denominator with a recorded discussion at their 16-week antenatal appointment of their preferred choice of birth setting.

Denominator – The number of women at low risk of complications attending a 16-week antenatal appointment.

Data source: Local data collection.

b) Proportion of women at low risk of complications with a recorded discussion at their 16-week antenatal appointment about local birth outcomes.

Numerator – The number in the denominator with a recorded discussion at their 16-week antenatal appointment about local birth outcomes.

Denominator – The number of women at low risk of complications attending a 16-week antenatal appointment.

Data source: Local data collection.

Outcome

Maternal experience and satisfaction with place of birth.

Data source: National data is collected as part of the [Care Quality Commission's Maternity Survey](#): question B3 ('Were you offered a choice about where to have your baby?').

What the quality statement means for different audiences

Service providers (community, primary and secondary care services) raise awareness of maternity pathways and ensure that systems and tools are in place to offer women at low risk of complications a choice of all 4 birth settings and local information about birth outcomes to support them to make informed decisions about where to have their baby.

Healthcare professionals provide women at low risk of complications with local information about birth outcomes and rates of transfer to an obstetric unit for all birth settings, and support them to make informed decisions about where to have their baby. Healthcare professionals can adapt and use [NICE's resource for midwives on choosing place of birth](#) to do this.

Commissioners (clinical commissioning groups or integrated care systems) commission maternity services to ensure that all 4 birth settings are available in the local or a neighbouring area to women at low risk of complications. Commissioners also ensure that services provide local information about outcomes for women and babies and rates of transfer to an obstetric unit for all birth settings to support women to make informed decisions about where to have their baby. Commissioners coordinate collection of outcome data in local and neighbouring areas to help service providers and healthcare professionals give information to women. Commissioners can refer to the [costing statement for the NICE guideline on intrapartum care for healthy women and babies](#) for more information about the likely resource impact of this quality statement, which will depend on local circumstances.

Women at low risk of having problems during labour and birth have a choice of 4 places where they can have their baby – at home, in a midwife-led unit that is either next to a hospital obstetric unit or in a different place, or in an obstetric unit ('labour ward'). To help women make an informed choice, they are given information by their midwife about birth outcomes and rates of transfer to an obstetric unit for their local or neighbouring area. Birth outcomes are things like the chances of needing a ventouse or forceps birth, caesarean birth or episiotomy, and the risk of serious medical problems for the baby.

Source guidance

- [Antenatal care. NICE guideline NG201](#) (2021), recommendation 1.3.13 and schedule of appointments (16-week appointment)
- [Intrapartum care for healthy women and babies. NICE guideline CG190](#) (2014, updated 2017), recommendation 1.1.2 (key priority for implementation), 1.1.3 and 1.1.6 (key priority for implementation)

Definitions of terms used in this quality statement

4 birth settings

The 4 settings where a woman at low risk of complications may choose to have her baby are: at home, in a freestanding midwifery unit, in an alongside midwifery unit and in an obstetric unit. [[NICE's guideline on intrapartum care for healthy women and babies](#), recommendation 1.1.2]

Birth outcomes

Outcomes for women for each planned place of birth include rates of spontaneous vaginal birth,

transfer to obstetric unit, obstetric intervention and delivering a baby with or without serious medical problems. [Adapted from [NICE's guideline on intrapartum care for healthy women and babies](#), recommendation 1.1.3]

Quality statement 2: One-to-one care

Quality statement

Women in established labour have one-to-one care and support from an assigned midwife.

Rationale

One-to-one care will increase the likelihood of the woman having a 'normal' vaginal birth without interventions, and will contribute to reducing both the length of labour and the number of operative deliveries. Care will not necessarily be given by the same midwife for the whole labour.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

Evidence of midwifery staff available to provide one-to-one care to women in established labour in each birth setting.

Data source: Local data collection.

Process

Midwifery staffing levels as in the [NICE guideline on safe midwifery staffing for maternity settings](#).

Numerator – The number of women in the denominator who receive one-to-one care from an assigned midwife during established labour.

Denominator – The number of women in established labour in a time period.

Data source: Local data collection.

Outcome

a) Neonatal morbidity.

Data source: Local data collection.

b) Maternal morbidity.

Data source: Local data collection. [MBRRACE-UK Confidential Enquiries into Maternal Deaths and Morbidity](#) and the [National Maternity and Perinatal Audit \(NMPA\) report on maternal morbidity](#). Indicators from MBRRACE-UK and the NMPA are available via the [National Maternity Services Dashboard](#).

c) Maternal satisfaction and experience of care.

Data source: National data is collected as part of the [Care Quality Commission's Maternity Survey](#): section C ('Your labour and the birth of your baby').

What the quality statement means for different audiences

Service providers (for all 4 birth settings) ensure that recommended midwifery staffing ratios are maintained so that women in established labour have one-to-one care and support from an assigned midwife.

Healthcare professionals (assigned midwives) give one-to-one care to each woman in established labour and are solely dedicated to the care of that woman.

Commissioners (clinical commissioning groups or integrated care systems) commission services that have systems in place to maintain recommended midwifery staffing ratios, so that women in established labour have one-to-one care and support from an assigned midwife. Commissioners can refer to the [costing statement for the NICE guideline on intrapartum care for healthy women and babies](#) for more information about the likely resource impact of this quality statement, which will depend on local circumstances.

A woman in labour is cared for by a midwife who is looking after just her – this is called 'one-to-one care'. She might not have the same midwife for the whole of labour. One-to-one care aims to ensure that the woman has a good experience of care and reduces the likelihood of problems for her and

her baby.

Source guidance

- [Intrapartum care for healthy women and babies. NICE guideline CG190 \(2014, updated 2017\)](#), recommendation 1.7.1 (key priority for implementation)
- [Safe midwifery staffing for maternity settings. NICE guideline NG4 \(2015\)](#), recommendation 1.2.2

Definitions of terms used in this quality statement

Established labour

Labour is established when:

- there are regular painful contractions and
- there is progressive cervical dilatation from 4 cm.

[[NICE's guideline on intrapartum care for healthy women and babies](#), recommendation 1.3.1]

Quality statement 3: Cardiotocography and initial assessment of a woman in labour

This statement has been removed. For more details see [update information](#).

Quality statement 4: Stopping cardiotocography

Quality statement

Women at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Rationale

Cardiotocography is offered to women if intermittent auscultation indicates possible fetal heart rate abnormalities. However, cardiotocography that is started for this reason should be stopped if the trace is normal for 20 minutes, because it restricts the woman's movement and can cause labour to slow down. This can lead to a cascade of interventions that may result in adverse birth outcomes.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

Evidence of local arrangements to ensure that women at low risk of complications having cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Data source: Local data collection.

Process

Proportion of women at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Numerator – The number in the denominator who have the cardiotocograph removed.

Denominator – The number of women in labour at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation and who have a normal trace for 20 minutes.

Data source: Local data collection.

Outcome

Maternal satisfaction and experience of care.

Data source: National data is collected as part of the [Care Quality Commission's Maternity Survey](#): section C ('Your labour and the birth of your baby').

What the quality statement means for different audiences

Service providers (for freestanding midwifery units, alongside midwifery units and obstetric units) have evidence of local arrangements to ensure that protocols are in place so that women in labour at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Healthcare professionals (midwives and obstetricians) remove the cardiotocograph if the trace is normal for 20 minutes for women at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation.

Commissioners (clinical commissioning groups or integrated care systems) specify and check that service providers have protocols in place to ensure that women in labour at low risk of complications who have cardiotocography because of concern arising from intermittent auscultation have the cardiotocograph removed if the trace is normal for 20 minutes.

Women who are at low risk of problems during labour, but who have electronic monitoring because of possible concerns about the baby's heartbeat, are taken off the monitor if the baby's heartbeat is normal for 20 minutes.

Source guidance

[Intrapartum care for healthy women and babies. NICE guideline CG190 \(2014, updated 2017\)](#), recommendations 1.4.12 and 1.10.8

Definitions of terms used in this quality statement

Normal cardiotocograph trace

A normal trace has the following normal/reassuring features:

- baseline fetal rate of 100 to 160 beats per minute and
- baseline variability of 5 to 25 beats per minute and
- no or early decelerations and
- variable decelerations with no concerning characteristics for less than 90 minutes.

It should be noted that while variable decelerations of less than 90 minutes is a reassuring feature, the trace should not be removed at 20 minutes if these are present without further assessment.

[[NICE's guideline on intrapartum care for healthy women and babies](#), table 10, and expert opinion]

Quality statement 5: Interventions during labour

Quality statement

Women at low risk of complications are not offered amniotomy or oxytocin if labour is progressing normally.

Rationale

For women at low risk of complications, amniotomy and oxytocin do not reduce the incidence of caesarean birth, increase the incidence of spontaneous vaginal births or contribute to improved neonatal outcomes. They are therefore unnecessary for women at low risk of complications if labour is progressing normally.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

Evidence of local arrangements to ensure that women at low risk of complications who are in labour that is progressing normally do not have amniotomy or oxytocin.

Data source: Local data collection.

Process

Proportion of women at low risk of complications whose labour is progressing normally who do not have amniotomy or oxytocin.

Numerator – The number in the denominator who do not have amniotomy or oxytocin.

Denominator – The number of women at low risk of complications whose labour is progressing normally.

Data source: Local data collection.

Outcome

a) The number of women in labour that is progressing normally having amniotomy or oxytocin.

Data source: Local data collection.

b) Maternal satisfaction and experience of care.

Data source: National data is collected as part of the [Care Quality Commission's Maternity Survey](#): section C ('Your labour and the birth of your baby').

What the quality statement means for different audiences

Service providers (for all 4 birth settings) have protocols in place to ensure that women at low risk of complications whose labour is progressing normally are not offered amniotomy or oxytocin.

Healthcare professionals (midwives and obstetricians) do not offer amniotomy or oxytocin to women at low risk of complications whose labour is progressing normally.

Commissioners (clinical commissioning groups or integrated care systems) specify and check that service providers have protocols in place to ensure that women at low risk of complications whose labour is progressing normally are not offered amniotomy or oxytocin.

Women who are at low risk of having problems and whose labour is progressing normally are not offered amniotomy (having their waters broken) or oxytocin (a medicine given through a drip that speeds up labour).

Source guidance

[Intrapartum care for healthy women and babies. NICE guideline CG190 \(2014, updated 2017\)](#), recommendations 1.12.11 and 1.12.12

Definitions of terms used in this quality statement

Normal labour and normal progression of labour

The [NICE full guideline on intrapartum care for healthy women and babies](#) adopts the World Health Organization definition of a normal labour: 'labour is normal when it is spontaneous in onset, low risk at the start and remaining so throughout labour and birth. The baby is born spontaneously and in the vertex position between 37 to 42 completed weeks of pregnancy. After birth woman and baby are in good condition'.

Quality statement 6: Delayed cord clamping

Quality statement

Women do not have the cord clamped earlier than 1 minute after the birth unless there is concern about cord integrity or the baby's heartbeat.

Rationale

The benefits of delayed cord clamping include higher haemoglobin concentrations, a decreased risk of iron deficiency and greater vascular stability in babies. If they wish, women can ask healthcare professionals to wait longer to clamp the cord.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

Evidence of local arrangements to ensure that midwives and obstetricians do not clamp the cord earlier than 1 minute after the birth unless there is a concern about cord integrity or the baby's heartbeat.

Data source: Local data collection.

Process

a) Proportion of cords clamped earlier than 1 minute after the birth where there is not a concern about cord integrity or the baby's heartbeat.

Numerator – The number in the denominator where the cord is clamped after 1 minute after the birth.

Denominator – The number of babies born where there is no concern about cord integrity or the baby's heartbeat.

Data source: Local data collection.

b) Proportion of cords clamped earlier than 1 minute where there is a concern about cord integrity or the baby's heartbeat.

Numerator – The number in the denominator where the cord is clamped earlier than 1 minute after the birth.

Denominator – the number of babies born where there is a concern about cord integrity or the baby's heartbeat.

Data source: Local data collection.

Outcome

Maternal satisfaction and experience of care.

Data source: National data is collected as part of the [Care Quality Commission's Maternity Survey: section C \('Your labour and the birth of your baby'\)](#).

What the quality statement means for different audiences

Service providers (for all 4 birth settings) have protocols in place to ensure that the cord is not clamped earlier than 1 minute after the birth unless there is concern about cord integrity or the baby's heartbeat.

Healthcare professionals (midwives and obstetricians) do not clamp the cord earlier than 1 minute after the birth unless there is concern about cord integrity or the baby's heartbeat.

Commissioners (clinical commissioning groups or integrated care systems) specify and check that service providers have protocols in place to ensure that the cord is not clamped earlier than 1 minute after the birth unless there is a concern about cord integrity or the baby's heartbeat.

Women who have just given birth do not have the cord clamped for at least 1 minute after the birth unless there are concerns about the baby. This is to allow more blood to reach the baby and may help to prevent anaemia.

Source guidance

Intrapartum care for healthy women and babies. NICE guideline CG190 (2014, updated 2017), recommendation 1.14.14 (key priority for implementation)

Definitions of terms used in this quality statement

Cord integrity

Concerns would arise over cord integrity if the cord was damaged in any way, if it had snapped during delivery or if there was bleeding to the cord. Definitions of cord integrity are not limited to those stated here. [Expert opinion]

Concern about the baby's heartbeat

Concern would arise if, after delivery, the baby has a heartbeat below 60 beats/minute that is not getting faster. [Adapted from NICE's guideline on intrapartum care for healthy women and babies, recommendation 1.14.14]

Quality statement 7: Skin-to-skin contact

Quality statement

Women have skin-to-skin contact with their babies after the birth.

Rationale

Skin-to-skin contact with babies soon after birth has been shown to promote the initiation of breastfeeding and protect against the negative effects of mother–baby separation.

Quality measures

The following measures can be used to assess the quality of care or service provision specified in the statement. They are examples of how the statement can be measured and can be adapted and used flexibly.

Structure

Evidence of local arrangements to ensure that midwives and obstetricians encourage women to have skin-to-skin contact with their babies after the birth.

Data source: Local data collection.

Process

Proportion of women with a record of having skin-to-skin contact with their babies after the birth. (Note: It is important that this happens as soon as possible, but timescales should be determined locally, depending on the setting and whether the baby and mother are stable.)

Numerator – The number in the denominator where there is a record of the woman having skin-to-skin contact with the baby.

Denominator – The number of babies born.

Data source: Local data collection. The [Maternity Services Data Set](#) collects data on whether a mother had skin-to-skin contact with their baby within 1 hour of birth.

Outcome

Women's satisfaction with the support received to have skin-to-skin contact with their babies after the birth.

Data source: Local data collection.

What the quality statement means for different audiences

Service providers (for all 4 birth settings) have protocols in place for midwives and obstetricians to encourage women to have skin-to-skin contact with their babies as soon as possible after the birth.

Healthcare professionals (midwives and obstetricians) encourage women to have skin-to-skin contact with their babies as soon as possible after the birth.

Commissioners (clinical commissioning groups or integrated care systems) specify and check that service providers have protocols in place to ensure that women are encouraged to have skin-to-skin contact with their babies as soon as possible after the birth.

Women are encouraged to have skin-to-skin contact with their babies as soon as possible after the birth.

Source guidance

[Intrapartum care for healthy women and babies. NICE guideline CG190 \(2014, updated 2017\), recommendation 1.15.6 \(key priority for implementation\)](#)

Update information

February 2017: Statement 3 on cardiotocography and the initial assessment of a woman in labour has been removed. This change has been made because the source guidance for this statement ([NICE's guideline on intrapartum care for healthy women and babies](#)) has been updated and the advice on cardiotocography for low risk women has changed.

Minor changes since publication

August 2021: Changes have been made to align statement 1 of this quality standard with [NICE's guideline on antenatal care](#). The process measures, recommendation numbers, references and links to the source guidance have been updated. Data sources and references have also been updated throughout.

November 2017: Changes were made to the definition of normal cardiotocograph trace in statement 4 to ensure alignment with [NICE's guideline on intrapartum care for healthy women and babies](#).

About this quality standard

NICE quality standards describe high-priority areas for quality improvement in a defined care or service area. Each standard consists of a prioritised set of specific, concise and measurable statements. NICE quality standards draw on existing NICE or NICE-accredited guidance that provides an underpinning, comprehensive set of recommendations, and are designed to support the measurement of improvement.

Expected levels of achievement for quality measures are not specified. Quality standards are intended to drive up the quality of care, and so achievement levels of 100% should be aspired to (or 0% if the quality statement states that something should not be done). However, this may not always be appropriate in practice. Taking account of safety, shared decision-making, choice and professional judgement, desired levels of achievement should be defined locally.

Information about [how NICE quality standards are developed](#) is available from the NICE website.

See our [webpage on quality standard advisory committees](#) for details of standing committee 2 members who advised on this quality standard. Information about the topic experts invited to join the standing members is available from the [webpage for this quality standard](#).

This quality standard has been included in the [NICE Pathways on safe midwifery staffing for maternity settings, intrapartum care and antenatal care](#), which bring together everything we have said on a topic in an interactive flowchart.

NICE has produced a [quality standard service improvement template](#) to help providers make an initial assessment of their service compared with a selection of quality statements. This tool is updated monthly to include new quality standards.

NICE guidance and quality standards apply in England and Wales. Decisions on how they apply in Scotland and Northern Ireland are made by the Scottish government and Northern Ireland Executive. NICE quality standards may include references to organisations or people responsible for commissioning or providing care that may be relevant only to England.

Resource impact

NICE quality standards should be achievable by local services. The potential resource impact is considered by the quality standards advisory committee, drawing on resource impact work for the

source guidance. Organisations are encouraged to use the resource impact products for the source guidance to help estimate local costs:

- [resource impact statement for NICE's guideline on antenatal care](#)
- [costing statement for NICE's guideline on intrapartum care for healthy women and babies.](#)

Diversity, equality and language

Equality issues were considered during development and [equality assessments for this quality standard](#) are available. Any specific issues identified during development of the quality statements are highlighted in each statement.

Commissioners and providers should aim to achieve the quality standard in their local context, in light of their duties to have due regard to the need to eliminate unlawful discrimination, advance equality of opportunity and foster good relations. Nothing in this quality standard should be interpreted in a way that would be inconsistent with compliance with those duties.

ISBN: 978-1-4731-1580-4

Endorsing organisation

This quality standard has been endorsed by NHS England, as required by the Health and Social Care Act (2012)

Supporting organisations

Many organisations share NICE's commitment to quality improvement using evidence-based guidance. The following supporting organisations have recognised the benefit of the quality standard in improving care for patients, carers, service users and members of the public. They have agreed to work with NICE to ensure that those commissioning or providing services are made aware of and encouraged to use the quality standard.

- [National Childbirth Trust](#)
- [Royal College of General Practitioners \(RCGP\)](#)
- [Royal College of Midwives](#)
- [Royal College of Obstetricians and Gynaecologists](#)