




Infigratinib

Revised: January 15, 2024.

CASRN: 872511-34-7



Drug Levels and Effects

Summary of Use during Lactation

Infigratinib is no longer marketed in the US. No information is available on the clinical use of infigratinib during breastfeeding. Because infigratinib is 96.8% bound to plasma proteins, the amount in milk is likely to be low. However, because of its potential toxicity in the breastfed infant and its half-life of 33.5 hours, the manufacturer recommends that breastfeeding be discontinued during infigratinib therapy and for 1 month after the last dose.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Infigratinib

CAS Registry Number

872511-34-7

Drug Class

Breast Feeding

Lactation

Milk, Human

Antineoplastic Agents

Enzyme Inhibitors

Protein Kinase Inhibitors

Receptor, Fibroblast Growth Factor, Type 1