




## Micafungin

Revised: May 15, 2023.

CASRN: 235114-32-6



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of micafungin during breastfeeding. Because micafungin is >99% bound to plasma proteins and has poor oral bioavailability, it is unlikely to reach the milk and be absorbed by the infant. Micafungin can safely be given intravenously to infants under 4 months of age. Any amount absorbed from milk is likely to be far less than an infant dose. If micafungin is required by the mother, it is not a reason to discontinue breastfeeding.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

*Infant Levels.* Relevant published information was not found as of the revision date.

### **Effects in Breastfed Infants**

Relevant published information was not found as of the revision date.

### **Effects on Lactation and Breastmilk**

Relevant published information was not found as of the revision date.

### **Alternate Drugs to Consider**

Fluconazole, Miconazole

## **Substance Identification**

### **Substance Name**

Micafungin

### **CAS Registry Number**

235114-32-6

### **Drug Class**

Breast Feeding

Lactation

Milk, Human

Antifungal Agents

Echinocandins