




## Vonoprazan

Revised: June 20, 2022.

CASRN: 881681-00-1



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the clinical use of vonoprazan during breastfeeding. Because of liver damage that occurred in nursing rodents, the manufacturer recommends that nursing mothers should pump and discard human milk while taking and for 2 days after the last dose. An alternate drug may be preferred.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Omeprazole, Pantoprazole,

## Substance Identification

### Substance Name

Vonoprazan

### CAS Registry Number

881681-00-1

### Drug Class

Breast Feeding

Lactation

Milk, Human

Anti-Ulcer Agents

Gastrointestinal Agents

Proton Pump Inhibitors