

**NLM Citation:** Drugs and Lactation Database (LactMed®) [Internet]. Bethesda (MD): National Institute of Child Health and Human Development; 2006-. Tremelimumab. [Updated 2023 Nov 15]. **Bookshelf URL:** https://www.ncbi.nlm.nih.gov/books/



# **Tremelimumab**

Revised: November 15, 2023.

CASRN: 745013-59-6

# **Drug Levels and Effects**

# **Summary of Use during Lactation**

No information is available on the clinical use of tremelimumab during breastfeeding. Because tremelimumab is a large protein molecule with a molecular weight of 149,000 Da, the amount in milk is likely to be very low.[1] It is also likely to be partially destroyed in the infant's gastrointestinal tract and absorption by the infant is probably minimal.[2] Until more data become available, toripalimab should be used with caution or avoided during breastfeeding, especially while nursing a newborn or preterm infant. Waiting for at least 2 weeks postpartum to resume therapy may minimize transfer to the infant.[3] The manufacturer recommends that women should not breastfeed during treatment and for 3 months after the last dose.

# **Drug Levels**

Maternal Levels. Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

#### **Effects in Breastfed Infants**

Relevant published information was not found as of the revision date.

#### **Effects on Lactation and Breastmilk**

Relevant published information was not found as of the revision date.

### References

- 1. Stratigakis A, Paty D, Zou P, et al. A regression approach for assessing large molecular drug concentration in breast milk. Reprod Breed 2023;3:199-207. doi:10.1016/j.repbre.2023.10.003
- 2. Anderson PO. Monoclonal antibodies during breastfeeding. Breastfeed Med 2021;16:591-3. PubMed PMID: 33956488.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

3. Krysko KM, Dobson R, Alroughani R, et al. Family planning considerations in people with multiple sclerosis. Lancet Neurol 2023;22:350-66. PubMed PMID: 36931808.

# **Substance Identification**

### **Substance Name**

Tremelimumab

# **CAS Registry Number**

745013-59-6

# **Drug Class**

**Breast Feeding** 

Lactation

Milk, Human

Antineoplastic Agents

Antibodies, Monoclonal, Humanized