




## Olutasidenib

Revised: January 15, 2023.

CASRN: 1887014-12-1



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of olutasidenib during breastfeeding. Because it is 93% bound to plasma proteins, the amount in milk is likely to be low. However, The manufacturer recommends that the mother not breastfeed during treatment and for 2 weeks after the last dose.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Olutasidenib

### CAS Registry Number

1887014-12-1

### Drug Class

Breast Feeding

Lactation

Milk, Human

Antineoplastic Agents

Enzyme Inhibitors

Isocitrate dehydrogenase-1 inhibitors