

## APPENDIX B. REFERENCES

- ACGIH (American Conference of Governmental Industrial Hygienists). (2021). 2021 TLVs and BEIs: Based on the documentation of the threshold limit values for chemical substances and physical agents & biological exposure indices. Cincinnati, OH.  
<https://portal.acgih.org/s/store#/store/browse/detail/a154W00000BOag7QAD>
- Anderson, D; Styles, JA. (1978). Appendix II. The bacterial mutation test: Six tests for carcinogenicity. Br J Cancer 37: 924-930. <http://dx.doi.org/10.1038/bjc.1978.134>
- Anderson, JG; Mccalla, DR; Bryant, DW; Mccarry, BE; Bromke, AM. (1987). Metabolic activation of 3-nitroperylene in the Salmonella/S9 assay. Mutagenesis 279: 279-285.  
<http://dx.doi.org/10.1093/mutage/2.4.279>
- Anderson, RS; Anderson, LM. (1987). Lack of effect of perylene as an initiator of lung tumors or as a promoter of skin tumors in mice [Abstract]. Fed Proc 46: 746.
- Archibong, AE; Inyang, F; Ramesh, A; Greenwood, M; Nayyar, T; Kopsombut, P; Hood, DB; Nyanda, AM. (2002). Alteration of pregnancy related hormones and fetal survival in F-344 rats exposed by inhalation to benzo(a)pyrene. Reprod Toxicol 16: 801-808.  
[http://dx.doi.org/10.1016/S0890-6238\(02\)00058-8](http://dx.doi.org/10.1016/S0890-6238(02)00058-8)
- Asada, S; Sasaki, K; Tanaka, N; Takeda, K; Hayashi, M; Umeda, M. (2005). Detection of initiating as well as promoting activity of chemicals by a novel cell transformation assay using v-Ha-ras-transfected BALB/c 3T3 cells (Bhas 42 cells). Mutat Res Genet Toxicol Environ Mutagen 588: 7-21. <http://dx.doi.org/10.1016/j.mrgentox.2005.07.011>
- Asokan, P; Das, M; Bik, DP; Howard, PC; Mccoy, GD; Rosenkranz, HS; Bickers, DR; Mukhtar, H. (1986). Comparative effects of topically applied nitrated arenes and their nonnitrated parent arenes on cutaneous and hepatic drug and carcinogen metabolism in neonatal rats. Toxicol Appl Pharmacol 86: 33-43. [http://dx.doi.org/10.1016/0041-008X\(86\)90397-2](http://dx.doi.org/10.1016/0041-008X(86)90397-2)
- ATSDR (Agency for Toxic Substances and Disease Registry). (1995). Toxicological profile for polycyclic aromatic hydrocarbons - Update [ATSDR Tox Profile]. (CIS/97/00215). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.  
<https://www.atsdr.cdc.gov/substances/toxsubstance.asp?toxicid=25>
- ATSDR (Agency for Toxic Substances and Disease Registry). (2021). Toxic substances portal: Toxicological profiles [Database]. Atlanta, GA. Retrieved from  
<https://www.atsdr.cdc.gov/toxprofiledocs/index.html>
- Bahri, R; Saidane-Mosbahi, D; Rouabha, M. (2010). Cytokine release and cytotoxicity in human keratinocytes induced by polycyclic aromatic hydrocarbons (1-methylpyrene and perylene). J Toxicol Environ Health A 73: 552-564.  
<http://dx.doi.org/10.1080/15287390903566617>
- Boyland, E; Sims, P. (1964). Metabolism of polycyclic compounds. 23. The metabolism of pyrene in rats and rabbits. Biochem J 90: 391-398. <http://dx.doi.org/10.1042/bj0900391>
- CalEPA (California Environmental Protection Agency). (2020). Consolidated table of OEHHA/CARB approved risk assessment health values. Sacramento, California.  
<https://ww2.arb.ca.gov/resources/documents/consolidated-table-oehha-carb-approved-risk-assessment-health-values>
- CalEPA (California Environmental Protection Agency). (2022). OEHHA chemical database [Database]. Sacramento, CA: Office of Environmental Health Hazard Assessment. Retrieved from <https://oehha.ca.gov/chemicals>
- Carver, JH; Machado, ML; MacGregor, JA. (1985). Petroleum distillates suppress in vitro metabolic activation: Higher [S-9] required in the Salmonella/microsome mutagenicity assay. Environ Mutagen 7: 369-379. <http://dx.doi.org/10.1002/em.2860070311>

- Carver, JH; Machado, ML; Macgregor, JA. (1986). Application of modified Salmonella/microsome prescreen to petroleum-derived complex mixtures and polynuclear aromatic hydrocarbons (PAH). *Mutat Res* 247: 247-253.  
[http://dx.doi.org/10.1016/0165-7992\(86\)90042-4](http://dx.doi.org/10.1016/0165-7992(86)90042-4)
- Casto, BC; Pieczynski, WJ; DiPaolo, JA. (1973). Enhancement of adenovirus transformation by pretreatment of hamster cells with carcinogenic polycyclic hydrocarbons. *Cancer Res* 33: 819-824.
- Casto, BC; Pieczynski, WJ; Janosko, N; DiPaolo, JA. (1976). Significance of treatment interval and DNA repair in the enhancement of viral transformation by chemical carcinogens and mutagens. *Chem Biol Interact* 13: 105-125.
- Chen, C; Tang, Y; Jiang, X; Qi, Y; Cheng, S; Qiu, C; Peng, B; Tu, B. (2012). Early postnatal benzo(a)pyrene exposure in Sprague-Dawley rats causes persistent neurobehavioral impairments that emerge postnatally and continue into adolescence and adulthood. *Toxicol Sci* 125: 248-261. <http://dx.doi.org/10.1093/toxsci/kfr265>
- Crespi, CL; Thilly, WG. (1984). Assay for gene mutation in a human lymphoblast line, AHH-1, competent for xenobiotic metabolism. *Mutat Res* 128: 221-230.  
[http://dx.doi.org/10.1016/0027-5107\(84\)90110-6](http://dx.doi.org/10.1016/0027-5107(84)90110-6)
- CTD (Comparative Toxicogenomics Database). (2022). Perylene: Comparative toxicogenomics database [Database]: MDI Biological Laboratory. Retrieved from <http://ctdbase.org/detail.go?type=chem&acc=D010569>
- Davis, AP; Grondin, CJ; Johnson, RJ; Sciaky, D; Wiegers, J; Wiegers, TC; Mattingly, CJ. (2021). The comparative toxicogenomics database (CTD): update 2021 [Database]: MDI Biological Laboratory. NC State University. Retrieved from <http://ctdbase.org/>
- De Flora, S. (1981). Study of 106 organic and inorganic compounds in the Salmonella/microsome test. *Carcinogenesis* 2: 283-298.  
<http://dx.doi.org/10.1093/carcin/2.4.283>
- De Flora, S; Zanacchi, P; Camoirano, A; Bennicelli, C; Badolati, GS. (1984). Genotoxic activity and potency of 135 compounds in the Ames reversion test and in a bacterial DNA-repair test [Review]. *Mutat Res* 133: 161-198. [http://dx.doi.org/10.1016/0165-1110\(84\)90016-2](http://dx.doi.org/10.1016/0165-1110(84)90016-2)
- DiPaolo, JA; Popescu, NC. (1974). Chromosome bands induced in human and syrian hamster cells by chemical carcinogens. *Br J Cancer* 30: 103-108.  
<http://dx.doi.org/10.1038/bjc.1974.120>
- Durant, JL; Busby, WF, Jr; Lafleur, AL; Penman, BW; Crespi, CL. (1996). Human cell mutagenicity of oxygenated, nitrated and unsubstituted polycyclic aromatic hydrocarbons associated with urban aerosols. *Mutat Res DNA Repair* 371: 123-157.  
[http://dx.doi.org/10.1016/S0165-1218\(96\)90103-2](http://dx.doi.org/10.1016/S0165-1218(96)90103-2)
- ECHA (European Chemicals Agency). (2022). Substance infocard: Perylene. Available online at <https://www.echa.europa.eu/web/guest/substance-information-/substanceinfo/100.005.365> (accessed April 5, 2022).
- El-Bayoumy, K; Hecht, SS; Hoffmann, D. (1982). Comparative tumor initiating activity on mouse skin of 6-nitrobenzo[a]pyrene, 6-nitrochrysene, 3-nitroperylene, 1-nitropyrene and their parent hydrocarbons. *Cancer Lett* 16: 333-337. [http://dx.doi.org/10.1016/0304-3835\(82\)90015-5](http://dx.doi.org/10.1016/0304-3835(82)90015-5)
- Finzi, C; Daudel, P; Prodi, G. (1968). Interference among polycyclic hydrocarbons in experimental skin carcinogenesis. *Eur J Cancer* 3: 497-501.  
[http://dx.doi.org/10.1016/0014-2964\(68\)90087-X](http://dx.doi.org/10.1016/0014-2964(68)90087-X)

- Flesher, JW; Myers, SR. (1990). Bioalkylation of benz(a)anthracene as a biochemical probe for carcinogenic activity: Lack of bioalkylation in a series of six noncarcinogenic polynuclear aromatic hydrocarbons. *Drug Metab Dispos* 18: 163-167.
- Florin, I; Rutberg, L; Curvall, M; Enzell, CR. (1980). Screening of tobacco smoke constituents for mutagenicity using the Ames' test. *Toxicology* 15: 219-232.  
[http://dx.doi.org/10.1016/0300-483X\(80\)90055-4](http://dx.doi.org/10.1016/0300-483X(80)90055-4)
- Goldfarb, JL; Suuberg, EM. (2008). Vapor pressures and enthalpies of sublimation of ten polycyclic aromatic hydrocarbons determined via the Knudsen effusion method. *Journal of Chemical and Engineering Data* 53: 670-676. <http://dx.doi.org/10.1021/je7005133>
- Gupta, RC; Earley, K; Sharma, S. (1988). Use of human peripheral blood lymphocytes to measure DNA binding capacity of chemical carcinogens. *PNAS* 85: 3513-3517.  
<http://dx.doi.org/10.1073/pnas.85.10.3513>
- Harris, M; Kamps, C; Safe, S. (1988). Role of the 4-5S binding protein in the induction of aryl hydrocarbon hydroxylase in the rat. *Carcinogenesis* 9: 1475-1480.  
<http://dx.doi.org/10.1093/carcin/9.8.1475>
- Hera, C; Pueyo, C. (1988). Response of the L-arabinose forward mutation assay of *Salmonella typhimurium* to frameshift-type mutagens. *Mutat Res* 203: 39-45.  
[http://dx.doi.org/10.1016/0165-1161\(88\)90006-4](http://dx.doi.org/10.1016/0165-1161(88)90006-4)
- Ho, CH; Clark, BR; Guerin, MR; Barkenbus, BD; Rao, TK; Epler, JL. (1981). Analytical and biological analyses of test materials from the synthetic fuel technologies: IV. Studies of chemical structure-mutagenic activity relationships of aromatic nitrogen compounds relevant to synfuels. *Mutat Res* 85: 335-345. [http://dx.doi.org/10.1016/0165-1161\(81\)90224-7](http://dx.doi.org/10.1016/0165-1161(81)90224-7)
- Horton, AW; Christian, GM. (1974). Cocarcinogenic versus incomplete carcinogenic activity among aromatic hydrocarbons: Contrast between chrysene and benzo(b)triphenylene. *J Natl Cancer Inst* 53: 1017-1020. <http://dx.doi.org/10.1093/jnci/53.4.1017>
- IARC (International Agency for Research on Cancer). (1983). Evaluation of the carcinogenic risk of chemicals to humans. International Agency for Research on Cancer. World Health Organization.
- IARC (International Agency for Research on Cancer). (1998). Perylene. In Polynuclear aromatic compounds, part 1: Chemical, environmental and experimental data Summary of data reported and evaluation (pp. 51). Lyon, France: World Health Organization.  
<https://monographs.iarc.fr/wp-content/uploads/2018/06/mono32.pdf>
- IARC (International Agency for Research on Cancer). (2010). Some non-heterocyclic polycyclic aromatic hydrocarbons and some related exposures [IARC Monograph]. Lyon, France.  
<http://monographs.iarc.fr/ENG/Monographs/vol92/mono92.pdf>
- IDEAconsult (IDEAconsult Ltd). (2018). Toxtree: Toxic hazard estimation by decision tree approach (Version 3.1.0) [Database]. Retrieved from  
<https://apps.ideaconsult.net/data/ui/toxtree>
- IPCS. (2020). INCHEM: Chemical safety information from intergovernmental organizations [Database]. Geneva, Switzerland: World Health Organization, Canadian Centre for Occupational Health and Safety. Inter-Organization Programme for the Sound Management of Chemicals. Retrieved from <http://www.inchem.org/>
- Kaden, DA; Hites, RA; Thilly, WG. (1979). Mutagenicity of soot and associated polycyclic aromatic hydrocarbons to *Salmonella typhimurium*. *Cancer Res* 39: 4152-4159.

- Lavoie, EJ; Bedenko, EV; Hirota, N; Hecht, SS; Hoffmann, D. (1979). A comparison of the mutagenicity, tumor-initiating activity and complete carcinogenicity of polynuclear aromatic hydrocarbons. In PW Jones; P Leber (Eds.), Polynuclear aromatic hydrocarbons (pp. 705-721). Ann Arbor, MI: Ann Arbor Science Publishers, Inc.
- Lipniak, M; Brandys, J. (1993). Toxicokinetics of fluoranthene, pyrene and benz(a)anthracene in the rat. *Polycycl Aromat Compd* 3: 111-119.  
<http://dx.doi.org/10.1080/10406639308047863>
- Lofroth, G; Toftgard, R; Nilsson, L; Agurell, E; J-A, G. (1984). Short-term bioassays of nitro derivatives of benzo(a)pyrene and perylene. *Carcinogenesis* 5: 925-930.  
<http://dx.doi.org/10.1093/carcin/5.7.925>
- Mersch-Sundermann, V; Mochayedi, S; Kekekordes, S. (1992). Genotoxicity of polycyclic aromatic hydrocarbons in Escherichia coli PQ37. *Mutat Res* 278: 1-9.  
[http://dx.doi.org/10.1016/0165-1218\(92\)90279-9](http://dx.doi.org/10.1016/0165-1218(92)90279-9)
- Mukhtar, H; Link, CM; Cherniack, E; Kushner, DM; Bickers, DR. (1982). Effect of topical application of defined constituents of coal tar on skin and liver aryl hydrocarbon hydroxylase and 7-ethoxycoumarin deethylase activities. *Toxicol Appl Pharmacol* 64: 541-549. [http://dx.doi.org/10.1016/0041-008X\(82\)90251-4](http://dx.doi.org/10.1016/0041-008X(82)90251-4)
- Neubert, D; Tapken, S. (1978). Some data on the induction of monooxygenases in fetal and neonatal mouse tissues. In Role of pharmacokinetics in prenatal and perinatal toxicology: Third Symposium on Prenatal Development, Berlin, 1978. Stuttgart, Germany: Thieme.
- Neubert, D; Tapken, S. (1988). Prenatal induction of benzo-a-pyrene hydroxylases in mice. *Arch Toxicol* 62: 192-199. <http://dx.doi.org/10.1007/bf00570139>
- NIOSH (National Institute for Occupational Safety and Health). (2018). NIOSH pocket guide to chemical hazards. Index of chemical abstracts service registry numbers (CAS No.). Atlanta, GA. <http://www.cdc.gov/niosh/npg/npgdcas.html>
- NIST (National Institute of Standards and Technology). (2022). NIST Chemistry WebBook, SRD 69. Perylene (CAS 198-55-0). Available online at <https://webbook.nist.gov/cgi/cbook.cgi?ID=C198550&Mask=4#ref-18>
- NLM (National Library of Medicine). (2022a). ChemIDplus advanced [Database]. Bethesda, MD: National Institutes of Health, National Library of Medicine. Retrieved from <https://chem.nlm.nih.gov/chemidplus/>
- NLM (National Library of Medicine). (2022b). PubChem compounds summary: Benzo[a]pyrene, 50-32-8. Available online at <https://pubchem.ncbi.nlm.nih.gov/compound/2336>
- NLM (National Library of Medicine). (2022c). PubChem compounds summary: Perylene, 198-55-0. Available online at <https://pubchem.ncbi.nlm.nih.gov/compound/9142>
- NLM (National Library of Medicine). (2022d). PubChem compounds summary: Pyrene, 129-00-0. Available online at <https://pubchem.ncbi.nlm.nih.gov/compound/31423>
- NTP (National Toxicology Program). (2021). Report on carcinogens, fifteenth edition. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service. <http://dx.doi.org/10.22427/NTP-OTHER-1003>
- OCHEM (Online Chemical Database). (2022). ToxAlerts (Version 4.2.151) [Database]: Online Chemical Database with Modeling Environment. Helmholtz Zentrum Muenchen - Deutsches Forschungszentrum fuer Gesundheit und Umwelt. Retrieved from <https://ochem.eu/alerts/home.do>
- OECD (Organisation for Economic Co-operation and Development). (2022). The OECD QSAR toolbox [version 4.5] [Database]. Retrieved from <http://www.oecd.org/chemicalsafety/risk-assessment/oecd-qsar-toolbox.htm>

- OSHA (Occupational Safety & Health Administration). (2021a). Air contaminants: Occupational safety and health standards for shipyard employment, subpart Z, toxic and hazardous substances. (OSHA Standard 1915.1000). Washington, DC. <https://www.osha.gov/laws-regs/regulations/standardnumber/1915/1915.1000>
- OSHA (Occupational Safety & Health Administration). (2021b). Safety and health regulations for construction: Occupational health and environmental controls: Gases, vapors, fumes, dusts, and mists. Washington, DC. <https://www.osha.gov/laws-regs/regulations/standardnumber/1926/1926.55>
- OSHA (Occupational Safety & Health Administration). (2021c). Table Z-1: Limits for air contaminants. Occupational safety and health standards, subpart Z, toxic and hazardous substances. Washington, DC. <https://www.osha.gov/laws-regs/regulations/standardnumber/1910/1910.1000TABLEZ1>
- Penman, BW; Kaden, DA; Liber, HL; Skopek, TR; Thilly, WG; Hites, RA. (1980). Perylene is a more potent mutagen than benzo(a)pyrene for *S. typhimurium*. *Mutat Res* 271: 271-277. [http://dx.doi.org/10.1016/0165-1218\(80\)90060-9](http://dx.doi.org/10.1016/0165-1218(80)90060-9)
- Piskorska-Pliszczynska, J; Keys, B; Safe, S; Newman, MS. (1986). The cytosolic receptor binding affinities and AHH induction potencies of 29 polynuclear aromatic hydrocarbons. *Toxicol Lett* 34: 67-74. [http://dx.doi.org/10.1016/0378-4274\(86\)90146-3](http://dx.doi.org/10.1016/0378-4274(86)90146-3)
- Popescu, NC; Amsbaugh, SC; DiPaolo, JA. (1980). Enhancement of N-methyl-N'-nitro-N-nitrosoguanidine transformation of Syrian hamster cells by a phorbol diester is independent of sister chromatid exchanges and chromosome aberrations. *Proc Natl Acad Sci USA* 7282: 7282-7286. <http://dx.doi.org/10.1073/pnas.77.12.7282>
- Popescu, NC; Turnbull, D; Dipaolo, JA. (1977). Sister chromatid exchange and chromosome aberration analysis with the use of several carcinogens and noncarcinogens. *J Natl Cancer Inst* 59: 289-293. <http://dx.doi.org/10.1093/jnci/59.1.289>
- Rády, P; Arany, I; Boján, F; Kertai, P. (1980). Effect of carcinogenic and non-carcinogenic chemicals on the activities of four glycolytic enzymes in mouse lung. *Chem Biol Interact* 31: 209-213. [http://dx.doi.org/10.1016/0009-2797\(80\)90007-1](http://dx.doi.org/10.1016/0009-2797(80)90007-1)
- Rády, P; Arany, I; Uzvolgyi, E; Bojan, F. (1981). Activity of pyruvate kinase and lactic acid dehydrogenase in mouse lung after transplacental exposure to carcinogenic and noncarcinogenic chemicals. *Toxicol Lett* 8: 223-228. [http://dx.doi.org/10.1016/0378-4274\(81\)90105-3](http://dx.doi.org/10.1016/0378-4274(81)90105-3)
- Reddy, MV; Gupta, RC; Randerath, E; Randerath, K. (1984). 32 P-postlabeling test for covalent DNA binding of chemicals in vivo: Application to a variety of aromatic carcinogens and methylating agents. *Carcinogenesis* 5: 231-244. <http://dx.doi.org/10.1093/carcin/5.2.231>
- Sakai, M; Yoshida, D; Mizusaki, S. (1985). Mutagenicity of polycyclic aromatic hydrocarbons and quinones on salmonella typhimurium TA97. *Mutat Res* 156: 61-67. [http://dx.doi.org/10.1016/0165-1218\(85\)90007-2](http://dx.doi.org/10.1016/0165-1218(85)90007-2)
- Salamone, MF; Haddle, JA; Katz, M. (1979). The mutagenic activity of thirty polycyclic aromatic hydrocarbons (PAH) and oxides in urban airborne particulates. *Environ Int* 2: 37-43. [http://dx.doi.org/10.1016/0160-4120\(79\)90092-8](http://dx.doi.org/10.1016/0160-4120(79)90092-8)
- Schwarzenbach, RP; Gschwend, PM; Imboden, DM. (2016). Environmental organic chemistry (3rd ed.). Hoboken, NJ: Wiley.
- Shimada, T; Fujii-Kuriyama, Y. (2004). Metabolic activation of polycyclic aromatic hydrocarbons to carcinogens by cytochromes P450 1A1 and 1B1 [Review]. *Cancer Sci* 95: 1-6. <http://dx.doi.org/10.1111/j.1349-7006.2004.tb03162.x>

- Sirianni, SR; Huang, CC. (1978). Sister chromatid exchange induced by promutagens/carcinogens in Chinese hamster cells cultured in diffusion chambers in mice. Proc Soc Exp Biol Med 158: 269-274. <http://dx.doi.org/10.3181/00379727-158-40186>
- TCEQ (Texas Commission on Environmental Quality). (2015). TCEQ guidelines to develop toxicity factors: revised September 2015. (RG-442). Austin, TX. <https://web.archive.org/web/20170305201245/http://www.tceq.texas.gov/publications/rgrg-442.html/>
- TCEQ (Texas Commission on Environmental Quality). (2022). TRRP protective concentration levels. Retrieved from <https://www.tceq.texas.gov/remediation/trrp/trppcls.html>
- Thayer, KA; Angrish, M; Arzuaga, X; Carlson, LM; Davis, A; Dishaw, L; Druwe, I; Gibbons, C; Glenn, B; Jones, R; Kaiser, JP; Keshava, C; Keshava, N; Kraft, A; Lizarraga, L; Persad, A; Radke, EG; Rice, G; Schulz, B; ... Vetter, N. (2022). Systematic evidence map (SEM) template: Report format and methods used for the US EPA Integrated Risk Information System (IRIS) program, Provisional Peer Reviewed Toxicity Value (PPRTV) program, and other "fit for purpose" literature-based human health analyses. Environ Int 169: 107468. <http://dx.doi.org/10.1016/j.envint.2022.107468>
- Thilly, WG; Longwell, J; Andon, BM. (1983). General approach to the biological analysis of complex mixtures. Environ Health Perspect 48: 129-136. <http://dx.doi.org/10.2307/3429464>
- U.S. EPA (U.S. Environmental Protection Agency). (1990). Integrated Risk Information System (IRIS) chemical assessment summary for pyrene (CASRN 129-00-0). Washington, DC: National Center for Environmental Assessment, Integrated Risk Information System. [https://cfpub.epa.gov/ncea/iris/iris\\_documents/documents/subst/0445\\_summary.pdf](https://cfpub.epa.gov/ncea/iris/iris_documents/documents/subst/0445_summary.pdf)
- U.S. EPA (U.S. Environmental Protection Agency). (1991). Guidelines for developmental toxicity risk assessment. Fed Reg 56: 63798-63826.
- U.S. EPA (U.S. Environmental Protection Agency). (1994). Methods for derivation of inhalation reference concentrations and application of inhalation dosimetry [EPA Report]. (EPA600890066F). Research Triangle Park, NC. <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=71993&CFID=51174829&CFTOKEN=25006317>
- U.S. EPA (U.S. Environmental Protection Agency). (2005). Guidelines for carcinogen risk assessment [EPA Report]. (EPA630P03001F). Washington, DC. [https://www.epa.gov/sites/production/files/2013-09/documents/cancer\\_guidelines\\_final\\_3-25-05.pdf](https://www.epa.gov/sites/production/files/2013-09/documents/cancer_guidelines_final_3-25-05.pdf)
- U.S. EPA (U.S. Environmental Protection Agency). (2007). Provisional peer-reviewed toxicity values for pyrene (CASRN 129-00-0) [EPA Report]. Cincinnati, OH. <https://cfpub.epa.gov/ncea/pprtv/documents/Pyrene.pdf>
- U.S. EPA (U.S. Environmental Protection Agency). (2011a). Chemical assessment clustering engine (ChemACE). Retrieved from <https://www.epa.gov/tsca-screening-tools/chemical-assessment-clustering-engine-chemace>
- U.S. EPA (U.S. Environmental Protection Agency). (2011b). Health effects assessment summary tables (HEAST) for superfund [EPA Report]. Washington, DC. <https://epa-heast.ornl.gov/heast.php>
- U.S. EPA (U.S. Environmental Protection Agency). (2012). Estimation Programs Interface Suite™ for Microsoft® Windows, v 4.11 [Computer Program]. Washington, DC. Retrieved from <https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface>

- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2017a). Toxicological review of benzo[a]pyrene (CASRN 50-32-8) [EPA Report]. (EPA635R17003Fa). Washington, DC: U.S. Environmental Protection Agency, Integrated Risk Information System.  
[https://cfpub.epa.gov/ncea/iris/iris\\_documents/documents/toxreviews/0136tr.pdf](https://cfpub.epa.gov/ncea/iris/iris_documents/documents/toxreviews/0136tr.pdf)
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2017b). Toxicological review of benzo[a]pyrene (CASRN 50-32-8): Supplemental information [EPA Report]. (EPA635R17003Fb). Washington, DC: U.S. Environmental Protection Agency, Integrated Risk Information System.  
[https://cfpub.epa.gov/si/si\\_public\\_file\\_download.cfm?p\\_download\\_id=535926&Lab=NC\\_EA](https://cfpub.epa.gov/si/si_public_file_download.cfm?p_download_id=535926&Lab=NC_EA)
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2018). 2018 Edition of the drinking water standards and health advisories tables [EPA Report]. (EPA 822-F-18-001). Washington, DC: U.S. Environmental Protection Agency, Office of Water.  
<https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=P100U7U8.txt>
- [U.S. EPA](#). (2020a). CompTox Chemicals Dashboard: Perylene [Bioactivity, invitrodb version 3.3] [Database]. Retrieved from  
<https://comptox.epa.gov/dashboard/chemical/details/DTXSID4047753>
- [U.S. EPA](#). (2020b). CompTox Chemicals Dashboard: Perylene [Bioactivity, PubChem] [Database]. Retrieved from  
<https://comptox.epa.gov/dashboard/chemical/details/DTXSID4047753>
- [U.S. EPA](#). (2020c). CompTox Chemicals Dashboard: Perylene [GenRA] [Database]. Retrieved from <https://comptox.epa.gov/dashboard/chemical/details/DTXSID4047753>
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2021). TSCA chemical substance inventory. Download the non-confidential TSCA inventory. Available online at  
<https://www.epa.gov/tsc-inventory/how-access-tsc-inventory>
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2022a). CompTox chemicals dashboard. Washington, DC. Retrieved from <https://comptox.epa.gov/dashboard>
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2022b). CompTox chemicals dashboard: Benzo(a)pyrene. Available online at  
<https://comptox.epa.gov/dashboard/chemical/details/DTXSID2020139> (accessed August 2, 2022).
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2022c). CompTox chemicals dashboard: Perylene [Database]. Retrieved from  
<https://comptox.epa.gov/dashboard/chemical/details/DTXSID4047753>
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2022d). CompTox chemicals dashboard: Pyrene. Available online at  
<https://comptox.epa.gov/dashboard/chemical/details/DTXSID3024289> (accessed August 2, 2022).
- [U.S. EPA](#) (U.S. Environmental Protection Agency). (2022e). Integrated risk information system (IRIS) database [Database]. Washington, DC. Retrieved from <http://www.epa.gov/iris/>
- [Van Duuren, BL; Sivak, A; Goldschmidt, BM; Katz, C; Melchionne, S.](#) (1970). Initiating activity of aromatic hydrocarbons in two-stage carcinogenesis. J Natl Cancer Inst 44: 1167-1173.
- [von der Hude, W; Behm, C; Gürtler, R; Basler, A.](#) (1988). Evaluation of the SOS chromotest. Mutat Res 203: 81-94. [http://dx.doi.org/10.1016/0165-1161\(88\)90023-4](http://dx.doi.org/10.1016/0165-1161(88)90023-4)
- [Wang, NC; Zhao, QJ; Wesselkamper, SC; Lambert, JC; Petersen, D; Hess-Wilson, JK.](#) (2012). Application of computational toxicological approaches in human health risk assessment. I. A tiered surrogate approach. Regul Toxicol Pharmacol 63: 10-19.  
<http://dx.doi.org/10.1016/j.yrtph.2012.02.006>

- White, KL; Lysy, HH; Holsapple, MP. (1985). Immunosuppression by polycyclic aromatic hydrocarbons: A structure-activity relationship in B6C3F1 and DBA/2 mice. *Immunopharmacology* 9: 155-164. [http://dx.doi.org/10.1016/0162-3109\(85\)90011-6](http://dx.doi.org/10.1016/0162-3109(85)90011-6)
- Zhao, W; Ramos, KS. (1995). Inhibition of DNA synthesis in primary cultures of adult rat hepatocytes by benzo[a]pyrene and related aromatic hydrocarbons: Role of Ah receptor-dependent events. *Toxicology* 99: 179-189. [http://dx.doi.org/10.1016/0300-483X\(94\)03028-Z](http://dx.doi.org/10.1016/0300-483X(94)03028-Z)