

Glycine Receptor Diversity in the Mammalian Retina

Silke Haverkamp, PhD

Created: March 27, 2012.

1 Introduction

Glycine is a major inhibitory neurotransmitter in the mammalian central nervous system (CNS). Its receptors, the inhibitory glycine receptors (GlyRs), are ligand-gated chloride channels composed of ligand-binding α and β subunits (1, 2). In mature neurons, the activation of GlyRs allows for an influx of chloride ions into the cytoplasm, which hyperpolarizes the postsynaptic membrane and, thereby, reduces neuronal firing. Inversely, the blockade of GlyRs by the competitive antagonist strychnine causes overexcitation resulting in pain, muscle cramps and exaggerated startle responses (3). Apart from its major transmitter function in spinal cord and brainstem, glycine also mediates substantial inhibitory neurotransmission via glycinergic amacrine cells in the mammalian retina. Thus, the abundant and highly complex expression patterns of different GlyR subtypes in the inner plexiform layer create an appealing field of retinal research.

2 Structure of glycine receptors

The GlyR was the first neurotransmitter receptor protein to be isolated from the mammalian CNS. Purification of the GlyR from rat spinal cord by strychnine affinity chromatography revealed three distinct polypeptides of molecular mass 48, 58 and 98 kDa (4). The 48 and 58 kDa peptides were shown to correspond to the α 1 and β subunits, respectively. The 98 kDa peptide was later identified as the cytoplasmic protein, gephyrin, which is essential for clustering GlyRs at postsynaptic densities via direct interactions between the GlyR β subunit and intracellular microtubules (5).

GlyRs are pentameric ligand-gated ion channel receptors of the Cys-loop family, which also include GABA_{A/C} receptors, muscle and neuronal nicotinic acetylcholine receptors, and serotonin type 3 receptors. Members of this superfamily share a common proposed structure (Fig. 1). Each subunit consists of a large N-terminal extracellular domain, four transmembrane segments (TM1–TM4), a long intracellular loop connecting TM3 and TM4, and a short extracellular C-terminus. The second transmembrane segment, TM2, lines the inner ion pore which, in GlyRs and GABA_{A/C} receptors, displays strict anion selectivity.

Author Affiliation: 1 Neuroanatomy, Max-Planck-Institute for Brain Research, Frankfurt/Main, Germany.; Email: silke.haverkamp@brain.mpg.de.

Corresponding author.

Copyright: © 2024 Webvision

Molecular cloning has revealed four genes encoding the α subunits ($\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 4$) and only one gene encoding the β subunit (6, 7). In the adult organism, two copies of the α subunit and three copies of the β subunit form the pentameric receptor protein (8).

3. Developmental expression

Glycine is excitatory during embryonic development and around birth. The neonatal form of the GlyR is thought to be a homopentamer of $\alpha 2$ subunits that are mainly found extrasynaptically in vivo, whereas adult synaptic GlyRs are heteromeric $\alpha\beta$ receptors. It seems that homomeric $\alpha 2$ GlyRs mediate a depolarizing glycine-gated chloride flux that in turn stimulates the calcium influx necessary for the development of numerous neuronal specializations, including glycinergic synapses (2, 9). Surprisingly, however, knockout of the $\alpha 2$ subunit has no obvious effect on neuronal development (10), whereas an $\alpha 1$ knockout (*Glra1^{spd-ot},oscillator*) has severe consequences: mice appear normal until the 2nd postnatal week whereupon they show prolonged periods of rapid tremor, producing extreme rigor and stiffness, and die within 10 days (11, 12).

4. Glycinergic amacrine cells

In the retina, approximately half of the amacrine cells release glycine at their synapses with bipolar, other amacrine, and ganglion cells. Glycinergic amacrine cells can be immunolabeled with antibodies against glycine or against the glycine transporter GlyT1 (13-15). Figure 2 shows a vertical section through a mouse retina that was double immunolabeled for glycine and for GlyT1 (16). Strong glycine immunoreactivity can be observed in amacrine cell bodies and their dendrites descending into the inner plexiform layer (IPL) or in dendrites from interplexiform amacrine cells ascending into the outer plexiform layer (OPL). Weak glycine expression is also found in putative ON-cone bipolar cells in the outer half of the inner nuclear layer (INL). The section was also immunolabeled for GlyT1 which labels all glycinergic amacrine cells but not bipolar cells. Bipolar cells do not express GlyT1 but they receive glycine by diffusion through electrical synapses (gap junctions) from glycinergic amacrine cells (17). In other parts of the CNS, GlyT1 has been localized to glial cells, while GlyT2 is now known to represent the presynaptic neuronal glycine transporter. Surprisingly, GlyT2 does not appear to be expressed in the mammalian retina (18).

5. Morphological types of glycinergic amacrine cells

Glycinergic amacrine cells are small-field amacrine cells with principally vertically oriented dendrites, and they comprise more than 10 different morphological types (13, 19). Most of them have small, diffuse dendritic trees and perform local circuit operations between the different sublayers of the IPL.

The most prominent and also most numerous glycinergic amacrine cell is the AII amacrine cell which transfers the light signal from rod bipolar cells into the cone pathway (20). In the inner IPL, AII cells receive direct glutamatergic input from rod bipolar cells, but they are also engaged with ON-cone bipolar cell axon terminals via electrical synapses (gap junctions). In the outer IPL, AII cell lobular dendrites provide glycinergic, chemical output synapses onto OFF-cone bipolar cell axon terminals and dendrites of OFF ganglion cells. Further glycinergic, small-field amacrine cells were identified in the cat retina by combined Golgi-staining and glycine uptake (21), and Menger et al. (13) identified at least 8 different glycinergic amacrine cells in the rat retina (Fig. 3).

Recently it became possible to study glycinergic amacrine cells in the retina of transgenic mice which express green fluorescent protein (GFP) under the control of the Thy1 promotor (Thy1-GFP-O) (22, 23). Three such cells are illustrated in figure 4, with double labeling for calcium binding protein 5 (CaBP5) in order to reveal the different sublaminae of the IPL and possible bipolar cell candidates as synaptic partners. The cells in figures 4A




Figure 1. Structure of ligand-gated ion channels. A. Each GlyR subunit consists of a large N-terminal extracellular domain, four transmembrane segments (TM1–TM4), a long intracellular loop connecting TM3 and TM4, and a short extracellular C-terminus. B. Proposed structure of a ligand-gated ion channel. TM2 forms the lining of the ion pore. C. Glycine receptors are pentamers constructed from α and β subunits in a ratio of $2\alpha:3\beta$. From Moss and Smart, 2001 (54).




Figure 2. Glycinergic amacrine cells of the mouse retina. A vertical section was double immunolabeled for glycine (red) and the glycine transporter GlyT1 (green). The arrows indicate an interplexiform process ascending to the OPL (OPL: outer plexiform layer; INL: inner nuclear layer; IPL: inner plexiform layer; GCL: ganglion cell layer). From Haverkamp and Wässle, 2000 (16).

and B have small, diffuse dendritic trees; the cell in figure 4C has a bistratified appearance and has been identified as A8 cell in cats and human retinas (24-26) (see also chapter on 'Roles of Amacrine Cells', Webvision).




Figure 3. Glycinergic amacrine cells of the rat retina. From Menger, Pow and Wässle, 1998 (13).




Figure 4. Glycinergic amacrine cells expressing GFP in the Thy1-GFP-O mouse. Sections are double labeled for bipolar cell marker CaBP5 (red). CaBP5 is expressed in rod bipolar cells (RB), in type 5 ON-cone bipolar cells (b5), and in type 3 OFF-cone bipolar cells (b3) (55). A2 and A4 amacrine cells are classified according to the scheme of Menger et al., 1998 (13); A8 according to Kolb et al., 1981 (26). Scale bar: 25 μ m. Unpublished data.

6. GlyR diversity in the mouse retina

The diversity of types of glycinergic amacrine cells is paralleled by the striking heterogeneity of glycine receptors. All four α subunits of the GlyR have been localized to specific synapses within the mammalian retina (23, 27-30). When subunit selective antibodies were applied to lightly fixed tissue, they each produced a distinct punctate immunofluorescence pattern (Figure 5).

The GlyRa1 subunit is expressed in a sparse population of puncta in the OPL, which represent synapses between glycinergic interplexiform cells and bipolar cell dendrites (Fig. 5A). In the outer IPL (stratum S1 and S2) GlyRa1 immunoreactivity is found in large puncta, which occur at high density. They represent synapses between AII amacrine cells and OFF-cone bipolar cells (30). In the inner IPL (stratum S3-S5) smaller GlyRa1 immunoreactive puncta can be observed representing synapses onto ganglion cell dendrites and rod bipolar cell axons (31, 32). The GlyRa2 subunit is more uniformly distributed across all strata (Fig. 5B), and GlyRa2 immunoreactive puncta occur at the highest density amongst the four α subunits (28). The GlyRa3 subunit (Fig. 5C) shows four bands whose density of puncta successively declines towards the ON sublamina (27). Lastly, the GlyRa4 subunit (Fig. 5D) shows a band of high density of puncta at the border between stratum 3 and 4 and further small and sparsely distributed puncta throughout the remaining strata (23). Taken together, the characteristic distribution of subunits across the IPL suggests that the GlyR subtypes are expressed at different synapses and are involved with different neuronal circuits.

Electron microscopy confirms the notion that the immunofluorescent puncta represent clusters of GlyRs at postsynaptic sites. Antibodies against the $\alpha 1$ subunit recognize extracellular epitopes of the receptor. Thus, staining from preembedding immunoelectron-microscopic experiments appears in the synaptic cleft of the synapse (Fig. 6A). Accordingly, postembedding immunoelectron microscopy showed that the $\alpha 3$ subunit was localized at the postsynaptic membrane (Fig. 6B).

7. Co-localization of GlyR subunits at postsynaptic sites

Since synaptic GlyRs are composed of 2α and 3β subunits (8) there is the possibility that two different α subunits co-exist in a single heteromeric GlyR. In addition, it is possible that two different GlyR subtypes, such as $\alpha 2\beta$ and $\alpha 3\beta$ GlyRs co-distribute at the same postsynaptic sites. In both cases, the immunoreactive hot spots should coincide. However, when retinal sections were double labeled for the GlyRa1 subunit and the other three GlyRa subunits, no statistically significant coincidence rate of immunoreactive puncta was observed. When retinal sections were double labeled for the GlyRa2 and $\alpha 3$ subunits a coincidence rate of 26.7% was found (28). In retinal sections double labeled for the GlyRa3 and $\alpha 4$ subunits no significant coincidence rate was found (23). In sections double labeled for the GlyRa4 and $\alpha 2$ subunits, 31.5% of the $\alpha 4$ immunoreactive clusters also contained the $\alpha 2$ subunit (Fig. 7). The results indicate that postsynaptic GlyR clusters usually contain only one type of α subunit. The exception is approximately one third of synapses immunoreactive for GlyR $\alpha 2$ which can also contain the $\alpha 3$ or the $\alpha 4$ subunits.

8. Expression of GlyRs by identified neurons

In order to reveal the involvement of selected GlyR subtypes with different retinal circuits, identified neurons were immunostained for the different GlyR α subunits. Figure 8 shows a GFP labeled A-type ganglion cell in a whole mount of Thy1-GFP-O mouse retina (32). The retina was immunolabeled for the GlyRa1 subunit demonstrating that many GlyRa1 immunoreactive puncta decorate the dendrites of this A-type ON ganglion cell. A coincidence of puncta and dendrites was also obvious for A-type OFF ganglion cells, which suggests that both ON and OFF A-type ganglion cells receive glycinergic input through $\alpha 1$ subunit expressing synapses. A-type ganglion cells were also double labeled for the other α subunits but quantification showed that the predominant input is through GlyRa1 containing synapses (32).

Figure 9 shows a vertical view of a Type 3 glycinergic amacrine cell in the Thy1-GFP-O mouse retina (33). Many GlyRa2 immunoreactive puncta (red) coincide with dendritic varicosities of the Type 3 cell (green) (Fig. 9D-F). Since this cell is a glycinergic amacrine cell, the puncta may represent input synapses the cell receives from other glycinergic amacrine cells or output synapses the cell makes onto other, non-labeled neurons. The fact that the red GlyRa2 immunoreactive puncta are always slightly displaced from the green varicosities (Fig. 9F) indicates that these synaptic GlyRa2 clusters are expressed by unknown neurons that are postsynaptic to this Type 3 cell.

The two examples of cells presented in figure 8 and figure 9 suggest a correlation between the morphological type of a given neuron and the molecular signature of the glycinergic synapse it receives or forms onto postsynaptic neurons. In this context one interesting question is whether the presynaptic neuron instructs the postsynaptic cell to express certain GlyR subunits or whether a given postsynaptic neuron expresses an exclusive GlyR subtype. To address this question a detailed physiological characterization of selected synaptic GlyRs on individual cells is essential.

To date, selective agonists or antagonists that distinguish different isoforms of synaptic GlyRs have not been identified (1, 34). However, mutant mice are available that have dysfunction of specific GlyR subunits and thus it became possible to study details of the glycinergic synaptic transmission in the mammalian retina (31, 32, 35, 36). The kinetic properties of GlyRs were measured by recording spontaneous inhibitory postsynaptic currents (sIPSCs) from identified retinal neurons in wild-type mice and mice lacking GlyR α subunit (*Gla1^{spd-ot}*, *Gla2*




Figure 5. Diversity of GlyR subtypes in the mouse retina. A. GlyR α 1 immunoreactive puncta are most prominent in the outer IPL (OFF sublamina). B. GlyR α 2 immunofluorescence is more evenly distributed across the IPL. C. GlyR α 3 expression is found in four bands. D. GlyR α 4 is most prominent in a small band at sublamina 3/4 border. Scale bar: 50 μ m. From Heinze et al., 2007 (23).




Figure 6. Synaptic localization of two GlyR subunits at conventional synapses in the IPL. A. Preembedding electron microscopy on vibratome sections of rat retina. The figure shows an amacrine cell synapse (AC, arrow) with GlyR α 1 immunoreactivity. From Sassoè-Pognetto et al., 1994 (30). B. Postembedding electron microscopy of GlyR α 3 on ultrathin sections of mouse retina. The GlyR subunit is localized at the postsynaptic membrane of an amacrine cell synapse (provided by Christian Puller, unpublished data).

and *Gla3*. From the observed differences of sIPSCs in wild-type and mutant mice, the cell-type specific subunit composition of GlyRs can be defined.

Glycine receptors expressed by bipolar cells

Patch-clamp recordings performed from identified bipolar cells in slices of the mouse retina (31) enabled the study of GlyRs by the application of exogenous glycine and by recording glycinergic spontaneous inhibitory




Figure 7. Colocalization of GlyR subunits at postsynaptic sites. GlyRa3 and GlyRa1 immunoreactive puncta are not colocalized whereas GlyRa2 and GlyRa4 immunoreactive puncta sometimes colocalize. From Haverkamp et al., 2003 (27); Heinze et al., 2007 (23).




Figure 8. Colocalization of GlyRa1 with the dendrites of an A-type ganglion cell. The image on the left shows an A2-ON ganglion cell in the Thy1-GFP-O mouse (22). The whole mount was also immunostained for GlyRa1 and many immunoreactive puncta decorate the dendrites of this ganglion cell. Scale bar: 20 μm . From Majumdar et al., 2007 (32).

postsynaptic currents (sIPSCs). Glycine application elicited large-amplitude currents in all OFF-cone and rod bipolar cells while ON-cone bipolar cells exhibited only very small, if any, glycinergic currents (Fig. 10 (31, 37),). By comparing sIPSCs of bipolar cells in wild-type and *Gla3*^{-/-} mice, no statistically significant differences were found; whereas glycine-induced currents and sIPSCs were absent from all bipolar cells of *Gla1*^{sp^d-ot} mice. Thus, OFF-cone and rod bipolar cells receive kinetically fast glycinergic inputs, preferentially mediated by GlyRs composed of $\alpha 1$ and β subunits (decay time constant $\tau \sim 5$ ms). Slow sIPSCs, the characteristic of GlyRs containing the $\alpha 2$ subunit, were not observed in bipolar cells.




Figure 9. Association of GlyRa2 with a glycinergic amacrine cell. C. Type 3 amacrine cell from the Thy1-GFP-O mouse retina. D. Single confocal section of the cell in C also immunostained for GlyRa2. The boxed area is shown at higher magnification in E and F. Scale bar: 17 μm in C & D, 10 μm in E & F. From Wässle et al., 2009 (33).




Figure 10. Glycine-induced currents from mouse bipolar cells. OFF- and ON-bipolar cells of the mouse retina as described by Ghosh et al., 2004 (55) and summary diagram of the peak amplitudes of inward currents elicited by the application of glycine (1–2 mM) in the nine cone bipolar cell types and rod bipolar (RB) cells of wild-type, *Glra3*^{-/-} and *Glra1*^{spd-ot} mice. From Ivanova et al., 2006 (31).

Glycine receptors on All and narrow-field amacrine cells

Amacrine cells are known to express strychnine-sensitive glycine receptors. The GlyRs expressed by All amacrine cells and by the narrow-field (NF) amacrine cells (Fig. 11A) were studied by patch-clamp recordings in mouse retinal slices (36). Glycinergic sIPSCs of All cells displayed medium fast kinetics ($\tau \sim 11$ ms, Fig. 11B), which were completely absent in the *Glra3*^{-/-} mouse, indicating that synaptic GlyRs of All cells mainly contain

the $\alpha 3$ subunit. Glycinergic sIPSCs of NF cells had slow kinetics ($\tau \sim 27$ ms, Figs. 11B and 12C) that were significantly prolonged in *Gla2*^{-/-} mice ($\tau \sim 69$ ms, Fig. 12F). These data show that morphologically distinct amacrine cells express different sets of glycine receptors.

Glycine receptors expressed by wide-field amacrine cells

Glycine induced currents and sIPSCs were also recorded from displaced wide-field, putative GABAergic amacrine cells. These GlyRs had slow kinetics ($\tau \sim 25$ ms) (35, 38) comparable to NF amacrine cells. ON-starburst (cholinergic amacrine cells) had sIPSCs with extremely long decay time constants ($\tau \sim 50 - 70$ ms), which did not differ between wild-type and the three mutant mice (35). Since GlyR $\alpha 4$ immunoreactive puncta (Fig. 5D) occur at higher density along the dendrites of ON-starburst amacrine cells, it is possible that GlyRs of ON-starburst cells are dominated by the $\alpha 4$ subunit. This would in turn suggest that GlyRs containing the $\alpha 4$ subunit have slow kinetics.

Glycine receptors expressed by ganglion cells

There are more than ten different types of ganglion cells in any mammalian retina. The GlyRs expressed by A-type ganglion cells (alpha-cell homologues) of the mouse retina were also investigated both in wild-type and mutant mice (32). In the wild-type retina, glycinergic sIPSCs of A-type ganglion cells have fast kinetics (mean $\tau = 3.9 \pm 2.5$ ms). Glycinergic sIPSCs recorded from *Gla2*^{-/-} and *Gla3*^{-/-} mice did not differ from those of wild-type mice. However, the number of glycinergic sIPSCs was significantly reduced in *Gla1*^{spd-ot} mice and the remaining sIPSCs had slower kinetics. These results show that A-type ganglion cells receive preferentially kinetically fast glycinergic inputs, mediated by GlyRs containing the $\alpha 1$ subunit.

B-type cells (beta-cell homologues) are probably involved with sustained neurotransmission. They are also believed to perform complex tasks like local edge detection. However, the specific roles of different B-type cells in the mouse retina are not yet clearly understood. There are four classes of small-field B-type ganglion cells (39), and there seems to be a substantial heterogeneity with respect to GlyR expression in these cell types. GlyRs of B1 cells are dominated by the $\alpha 1$ subunit, while in B4 cells the $\alpha 3$ subunit plays an important role. B2 and B3 cells express a more balanced mixture of fast ($\alpha 1$) and slow ($\alpha 2$, $\alpha 3$, $\alpha 4$) GlyR subunits (40).


9. Summary and conclusion

All four GlyR α subunits are clustered in synaptic hot spots (Fig. 5) that show characteristic distributions across the IPL of the mouse retina (23). Gephyrin is responsible for clustering GlyRs to postsynaptic sites by linking the GlyR β subunit to the cytoskeleton (5, 41). No GlyR clusters could be detected in gephyrin deficient mouse retinas (42), which suggests that synaptic GlyRs in the retina are always heteromeric. In the adult, two copies of the α subunit and three copies of the β subunit form the pentameric receptor protein (8).

Bipolar cells and A-type ganglion cells represent the fast transfer channel of the mammalian retina and, therefore, express $\alpha 1\beta$ GlyRs with fast kinetics. AII amacrine cells relay rod light signals with lower temporal resolution and express $\alpha 3\beta$ GlyRs with medium-fast kinetics. NF amacrine cells are modulatory interneurons, where temporal precision seems less important and they express the $\alpha 2\beta$ and $\alpha 4\beta$ GlyRs with slow kinetics.

Both GABA and glycine inhibition are used to modulate different aspects of ganglion cell responses and receptive field (RF) organization (43-45). GABAergic amacrine cells often provide feedforward inhibition and target both GABA_ARs and GABA_CRs in reciprocal inhibitory circuits to modulate the RF center excitation (46), and they target GABA_ARs to amplify and refine the RF surround (47). Glycinergic inhibition on the other hand modulates less spatial than temporal properties and may increase the gain of the ganglion cell response. Glycinergic amacrine cells often use (crossover) inhibition to modulate ganglion cell excitatory inputs within their RF center (48-52).

A



B




Figure 11. Glycinergic sIPSCs of mouse narrow-field amacrine cells. A. Examples of recorded All, and narrow field (NF) glycinergic amacrine cells. B. Frequency histogram of decay time constants of glycinergic sIPSCs recorded from All cells and from NF cells in

wild-type mice. From Weiss et al., 2008 (36).

About the Authors



Silke Haverkamp received her PhD from the Carl von Ossietzky Universität Oldenburg, Germany, in Visual Neuroscience under the mentorship of Josef Ammermüller. She was a postdoctoral researcher at the Boston University with Bill Eldred and then at the Moran Eye Center in Salt Lake City with Helga Kolb. Afterwards, Silke moved back to Germany to join the Department of Neuroanatomy headed by Heinz Wässle at the Max Planck Institute of Brain Research in Frankfurt. She has now her own research group at the Institute and focuses on the structure and function of neurons and synaptic circuits within the mammalian retina, using neuroanatomical and electrophysiological methods.




Figure 12. Mutations of glycine receptor α subunits change sIPSC decay times. C-F. Decay time constants of glycinergic sIPSCs

recorded from narrow field (NF) cells of wild-type, *Gla^{spd-ot}*, *Gla^{3-/-}* and *Gla^{2-/-}* mice. From Weiss et al., 2008 (36).

References

1. Betz H., Laube B. *Glycine receptors: recent insights into their structural organization and functional diversity.* . J Neurochem. 2006;97(6):1600–10. PubMed PMID: 16805771.
2. Lynch J.W. *Native glycine receptor subtypes and their physiological roles.* . Neuropharmacology. 2009;56(1):303–9. PubMed PMID: 18721822.
3. Harvey R.J.et al. *The genetics of hyperekplexia: more than startle!* . Trends Genet. 2008;24(9):439–47. PubMed PMID: 18707791.
4. Pfeiffer F., Graham D., Betz H. *Purification by affinity chromatography of the glycine receptor of rat spinal cord.* . J Biol Chem. 1982;257(16):9389–93. PubMed PMID: 6286620.
5. Meyer G.et al. *Identification of a gephyrin binding motif on the glycine receptor beta subunit.* . Neuron. 1995;15(3):563–72. PubMed PMID: 7546736.
6. Harvey R.J.et al. *Glycine receptors containing the alpha4 subunit in the embryonic sympathetic nervous system, spinal cord and male genital ridge.* . Eur J Neurosci. 2000;12(3):994–1001. PubMed PMID: 10762330.
7. Matzenbach B.et al. *Structural analysis of mouse glycine receptor alpha subunit genes. Identification and chromosomal localization of a novel variant.* . J Biol Chem. 1994;269(4):2607–12. PubMed PMID: 7507926.
8. Grudzinska J.et al. *The beta subunit determines the ligand binding properties of synaptic glycine receptors.* . Neuron. 2005;45(5):727–39. PubMed PMID: 15748848.
9. Kirsch J., Betz H. *Glycine-receptor activation is required for receptor clustering in spinal neurons.* . Nature. 1998;392(6677):717–20. PubMed PMID: 9565032.
10. Young-Pearse T.L.et al. *Characterization of mice with targeted deletion of glycine receptor alpha 2.* . Mol Cell Biol. 2006;26(15):5728–34. PubMed PMID: 16847326.
11. Buckwalter M.S.et al. *A frameshift mutation in the mouse alpha 1 glycine receptor gene (*Gla1*) results in progressive neurological symptoms and juvenile death.* . Hum Mol Genet. 1994;3(11):2025–30. PubMed PMID: 7874121.
12. Kling C.et al. *The frameshift mutation oscillator (*Gla1*(*spd-ot*)) produces a complete loss of glycine receptor alpha1-polypeptide in mouse central nervous system.* . Neuroscience. 1997;78(2):411–7. PubMed PMID: 9145798.
13. Menger N., Pow D.V., Wässle H. *Glycinergic amacrine cells of the rat retina.* . J Comp Neurol. 1998;401(1):34–46. PubMed PMID: 9802699.
14. Pow D.V. *Transport is the primary determinant of glycine content in retinal neurons.* . J Neurochem. 1998;70(6):2628–36. PubMed PMID: 9603230.
15. Pow D.V., Hendrickson A.E. *Distribution of the glycine transporter glyt-1 in mammalian and nonmammalian retinæ.* . Vis Neurosci. 1999;16(2):231–9. PubMed PMID: 10367958.
16. Haverkamp S., Wässle H. *Immunocytochemical analysis of the mouse retina.* . J Comp Neurol. 2000;424(1):1–23. PubMed PMID: 10888735.
17. Vaney D.I., Nelson J.C., Pow D.V. *Neurotransmitter coupling through gap junctions in the retina.* . J Neurosci. 1998;18(24):10594–602. PubMed PMID: 9852595.
18. Zafra F.et al. *Glycine transporters are differentially expressed among CNS cells.* . J Neurosci. 1995;15(5 Pt 2):3952–69. PubMed PMID: 7751957.
19. MacNeil M.A., Masland R.H. *Extreme diversity among amacrine cells: implications for function.* . Neuron. 1998;20(5):971–82. PubMed PMID: 9620701.
20. Kolb H., Famiglietti E.V. *Rod and cone pathways in the inner plexiform layer of cat retina.* . Science. 1974;186(4158):47–9. PubMed PMID: 4417736.
21. Pourcho R.G., Goebel D.J. *A combined Golgi and autoradiographic study of (³H)glycine-accumulating amacrine cells in the cat retina.* . J Comp Neurol. 1985;233(4):473–80. PubMed PMID: 2984258.

22. Feng G. et al. *Imaging neuronal subsets in transgenic mice expressing multiple spectral variants of GFP.* . Neuron. 2000;28(1):41–51. PubMed PMID: 11086982.
23. Heinze L. et al. *Diversity of glycine receptors in the mouse retina: localization of the alpha4 subunit.* . J Comp Neurol. 2007;500(4):693–707. PubMed PMID: 17154252.
24. Kolb H., Linberg K.A., Fisher S.K. *Neurons of the human retina: a Golgi study.* . J Comp Neurol. 1992;318(2):147–87. PubMed PMID: 1374766.
25. Kolb H., Nelson R. *Hyperpolarizing, small-field, amacrine cells in cone pathways of cat retina.* . J Comp Neurol. 1996;371(3):415–36. PubMed PMID: 8842896.
26. Kolb H., Nelson R., Mariani A. *Amacrine cells, bipolar cells and ganglion cells of the cat retina: a Golgi study.* . Vision Res. 1981;21(7):1081–1114. PubMed PMID: 7314489.
27. Haverkamp S. et al. *Diversity of glycine receptors in the mouse retina: localization of the alpha3 subunit.* . J Comp Neurol. 2003;465(4):524–39. PubMed PMID: 12975813.
28. Haverkamp S. et al. *Diversity of glycine receptors in the mouse retina: localization of the alpha2 subunit.* . J Comp Neurol. 2004;477(4):399–411. PubMed PMID: 15329889.
29. Jusuf P.R., Haverkamp S., Grunert U. *Localization of glycine receptor alpha subunits on bipolar and amacrine cells in primate retina.* . J Comp Neurol. 2005;488(2):113–28. PubMed PMID: 15924342.
30. Sassoe-Pognetto M., Wässle H., Grunert U. *Glycinergic synapses in the rod pathway of the rat retina: cone bipolar cells express the alpha 1 subunit of the glycine receptor.* . J Neurosci. 1994;14(8):5131–46. PubMed PMID: 8046473.
31. Ivanova E., Müller F. *Retinal bipolar cell types differ in their inventory of ion channels.* . Vis Neurosci. 2006;23(2):143–54. PubMed PMID: 16638168.
32. Majumdar S. et al. *Glycine receptors of A-type ganglion cells of the mouse retina.* . Vis Neurosci. 2007;24(4):471–87. PubMed PMID: 17550639.
33. Wässle H. et al. *Glycinergic transmission in the Mammalian retina.* . Front Mol Neurosci. 2009;2:6. PubMed PMID: 19924257.
34. Webb T.I., Lynch J.W. *Molecular pharmacology of the glycine receptor chloride channel.* . Curr Pharm Des. 2007;13(23):2350–67. PubMed PMID: 17692006.
35. Majumdar S., Weiss J., Wässle H. *Glycinergic input of widefield, displaced amacrine cells of the mouse retina.* . J Physiol. 2009;587(Pt 15):3831–49. PubMed PMID: 19528249.
36. Weiss J. et al. *Glycinergic input of small-field amacrine cells in the retinas of wildtype and glycine receptor deficient mice.* . Mol Cell Neurosci. 2008;37(1):40–55. PubMed PMID: 17920294.
37. Eggers E.D., Lukasiewicz P.D. *GABA_A, GABA_C and glycine receptor-mediated inhibition differentially affects light-evoked signalling from mouse retinal rod bipolar cells.* . J Physiol. 2006;572(Pt 1):215–25. PubMed PMID: 16439422.
38. Veruki M.L., Gill S.B., Hartveit E. *Spontaneous IPSCs and glycine receptors with slow kinetics in wide-field amacrine cells in the mature rat retina.* . J Physiol. 2007;581(Pt 1):203–19. PubMed PMID: 17331993.
39. Sun W., Li N., He S. *Large-scale morphological survey of mouse retinal ganglion cells.* . J Comp Neurol. 2002;451(2):115–26. PubMed PMID: 12209831.
40. Majumdar, S., L. Heinze, and H. Wässle, *Glycinergic Synaptic Inputs of B-type Ganglion Cells of the Mouse Retina.* ARVO Meeting Abstracts, 2007**48**(5): p. 4597.
41. Kirsch J., Betz H. *Widespread expression of gephyrin, a putative glycine receptor-tubulin linker protein, in rat brain.* . Brain Res. 1993;621(2):301–10. PubMed PMID: 8242343.
42. Fischer F. et al. *Reduced synaptic clustering of GABA and glycine receptors in the retina of the gephyrin null mutant mouse.* . J Comp Neurol. 2000;427(4):634–48. PubMed PMID: 11056469.
43. Caldwell J.H., Daw N.W. *Effects of picrotoxin and strychnine on rabbit retinal ganglion cells: changes in centre surround receptive fields.* . J Physiol. 1978;276:299–310. PubMed PMID: 650452.
44. Chen X. et al. *Three forms of spatial temporal feedforward inhibition are common to different ganglion cell types in rabbit retina.* . J Neurophysiol. 2010;103(5):2618–32. PubMed PMID: 20220071.
45. O'Brien B.J., Richardson R.C., Berson D.M. *Inhibitory network properties shaping the light evoked responses of cat alpha retinal ganglion cells.* . Vis Neurosci. 2003;20(4):351–61. PubMed PMID: 14658764.

46. Eggers E.D., Lukasiewicz P.D. *Multiple pathways of inhibition shape bipolar cell responses in the retina.* . Vis Neurosci. 2011;28(1):95–108. PubMed PMID: 20932357.
47. Flores-Herr N., Protti D.A., Wässle H. *Synaptic currents generating the inhibitory surround of ganglion cells in the mammalian retina.* . J Neurosci. 2001;21(13):4852–63. PubMed PMID: 11425912.
48. Hsueh H.A., Molnar A., Werblin F.S. *Amacrine-to-amacrine cell inhibition in the rabbit retina.* . J Neurophysiol. 2008;100(4):2077–88. PubMed PMID: 18667544.
49. Manookin M.B. et al. *Disinhibition combines with excitation to extend the operating range of the OFF visual pathway in daylight.* . J Neurosci. 2008;28(16):4136–50. PubMed PMID: 18417693.
50. van Wyk M., Wässle H., Taylor W.R. *Receptive field properties of ON- and OFF-ganglion cells in the mouse retina.* . Vis Neurosci. 2009;26(3):297–308. PubMed PMID: 19602302.
51. Wässle H., Schafer-Trenkler I., Voigt T. *Analysis of a glycinergic inhibitory pathway in the cat retina.* . J Neurosci. 1986;6(2):594–604. PubMed PMID: 3950712.
52. Werblin F.S. *Six different roles for crossover inhibition in the retina: correcting the nonlinearities of synaptic transmission.* . Vis Neurosci. 2010;27(1-2):1–8. PubMed PMID: 20392301.
53. Stafford B.K. et al. *Spatial-Temporal Patterns of Retinal Waves Underlying Activity-Dependent Refinement of Retinofugal Projections.* . Neuron. 2009;64(2):200–212. PubMed PMID: 19874788.
54. Moss S.J., Smart T.G. *Constructing inhibitory synapses.* . Nat Rev Neurosci. 2001;2(4):240–250. PubMed PMID: 11283747.
55. Ghosh K.K. et al. *Types of bipolar cells in the mouse retina.* . J Comp Neurol. 2004;469(1):70–82. PubMed PMID: 14689473.

License

All copyright for chapters belongs to the individual authors who created them. However, for non-commercial, academic purposes, images and content from the chapters portion of Webvision may be used with a non-exclusive rights under a Attribution, [Noncommercial 4.0 International \(CC BY-NC\) Creative Commons license](http://creativecommons.org/licenses/by-nc/4.0/). Cite Webvision, <http://webvision.med.utah.edu/> as the source. Commercial applications need to obtain license permission from the administrator of Webvision and are generally declined unless the copyright owner can/wants to donate or license material. Use online should be accompanied by a link back to the original source of the material. All imagery or content associated with blog posts belong to the authors of said posts, except where otherwise noted.